





# 11<sup>th</sup> ANNUAL HYBRID CONFERENCE

## American Council for Medicinally Active Plants (ACMAP)

JUNE 28 - JULY 2, 2022

Inter American University of Puerto Rico, Barranquitas Campus  
Puerto Rico, USA



## ABSTRACTS





**11<sup>th</sup> ANNUAL HYBRID CONFERENCE**  
**American Council for Medicinally Active Plants (ACMAP)**

**EXPLORING THE UNEXPLORED: SCIENCE  
AND APPLICATIONS OF MEDICINAL PLANTS**

**JUNE 28 - JULY 2, 2022 | PUERTO RICO, USA**

# 11<sup>th</sup> ANNUAL HYBRID CONFERENCE



## American Council for Medicinally Active Plants (ACMAP)

EXPLORING THE UNEXPLORED: SCIENCE AND APPLICATIONS OF MEDICINAL PLANTS

JUNE 28 - JULY 2, 2022 | PUERTO RICO, USA

Inter American University of Puerto Rico and its campuses in Barranquitas and Bayamón welcome you to the Eleventh Annual Conference of the American Council of Medicinally Active Plants (ACMAP). Both campuses are the hosts for this event and our entire university system feels extremely honored to receive and assist you in these days of arduous work.

This great scientific gathering that takes place from June 28th to July 2nd, 2022, is an important international event, with more than 60 scientists from the United States, Puerto Rico, Germany, India, Barbados, and Jamaica, including Dr. David Julius, Nobel Prize in Physiology in 2021, whom we salute and distinguish for joining us.

This Conference will allow for the discussion of various scientific topics, which will mostly revolve around the research, development, production, and conservation of plants with properties that help prevent and treat diseases.

This meeting facilitates direct contact and the exchange of ideas with professionals linked to the most recent discoveries in the realm of medicinally active plants. In addition, it represents an opportunity to promote faculty training and to stimulate an appreciation for research among our students.

We appreciate your participation and your commitment to science and research that benefits humanity. Welcome to the Inter American University of Puerto Rico.

Congratulations to the leadership of the Barranquitas and Bayamón Campuses



**DR. RAFAEL RAMÍREZ RIVERA**

INTERIM PRESIDENT OF THE  
INTER AMERICAN UNIVERSITY OF PUERTO RICO



# 11<sup>th</sup> ANNUAL HYBRID CONFERENCE



## American Council for Medicinally Active Plants (ACMAP)

EXPLORING THE UNEXPLORED: SCIENCE AND APPLICATIONS OF MEDICINAL PLANTS

JUNE 28 - JULY 2, 2022 | PUERTO RICO, USA

### HOSTS 2022

**DR. JUAN A. NEGRÓN BERRÍOS**  
INTER AMERICAN UNIVERSITY OF PUERTO RICO  
CHANCELLOR OF BARRANQUITAS CAMPUS

Inter American University of Puerto Rico at Barranquitas and Bayamón Campuses are pleased and honored to participate as the host institution for the 11th Annual hybrid conference of the American Council for Medically Active Plants (ACMAP). Traditional as well as modern medicine, have used plants products to preserve health and treat many diseases for humans and livestock. Plants offer an enormous amount of different chemicals compounds, with a wide range of applications, including medicinal properties. Plant biodiversity accounts for approximately 400,000 plant species, however, only a small number have been screened for medicinal properties. This implies a great opportunity for research and development, focusing in areas such as bioprospecting, clinical effects, mechanisms of action, structure-activity relationships, pharmacokinetic properties and many others. The celebration of Puerto Rico of the ACMAP 2022, offers the opportunity to examine the richness of medicinal plants locally, as well in the Caribbean. Medicinal plants in the Caribbean has a long cultural tradition, as well as scientifically. We see an opportunity for collaboration, in which bioprospecting of local flora for bioactive molecules for the identification of new leads and eventually into drugs.

I hope that all participants enjoy the conference and have a memorable experience. For those visiting, I wish a pleasant stay and the opportunity to appreciate rich cultural and social activities available in Puerto Rico.



**JUAN NEGRON-BERRIOS, PH.D.**

HOST UNIVERSITY CHANCELLOR



**CARLOS OLIVARES, PH.D.**

HOST UNIVERSITY CHANCELLOR



**ALOK ARUN, PH.D.**

CHAIR



**NIRMAL JOSHEE, PH.D.**

CO-CHAIR



# 11<sup>th</sup> ANNUAL HYBRID CONFERENCE



## American Council for Medicinally Active Plants (ACMAP)

EXPLORING THE UNEXPLORED: SCIENCE AND APPLICATIONS OF MEDICINAL PLANTS

JUNE 28 - JULY 2, 2022 | PUERTO RICO, USA

**DR. CARLOS OLIVARES**  
INTER AMERICAN UNIVERSITY OF PUERTO RICO  
CHANCELLOR OF BAYAMON CAMPUS

Welcome everyone to the Interamerican University of Puerto Rico, particularly to the Barranquitas and Bayamón campuses.

Medicinal plants have been used in healthcare since time immemorial. Studies have been carried out globally to verify their efficacy and some of the findings have led to the production of plant-based medicines.

Medicinal plants play vital roles in disease prevention and their promotion and use fit into all existing prevention strategies. However, conscious efforts need to be made to properly identify, recognize, and position medicinal plants in the design and implementation of these strategies. These approaches present interesting and emerging perspectives in the field of medicinal plants. Recommendations are proposed for strategizing the future role and place for medicinal plants in disease prevention.

This year's congress will focus on scientific advances in various areas of health and nutrition. We hope that this meeting will be a forum to reflect on research and development activities, both basic and applied, that demonstrate how the application of science can and does make a difference to communities and society.

I hope that this event is to the liking of all of you and that it serves to learn more about such an important topic as science and applications of medicinal plants.



### HOSTS 2022



**JUAN NEGRON-BERRIOS, PH.D.**

HOST UNIVERSITY CHANCELLOR



**CARLOS OLIVARES, PH.D.**

HOST UNIVERSITY CHANCELLOR



**ALOK ARUN, PH.D.**

CHAIR



**NIRMAL JOSHEE, PH.D.**

CO-CHAIR

# 11<sup>th</sup> ANNUAL HYBRID CONFERENCE



## American Council for Medicinally Active Plants (ACMAP)

EXPLORING THE UNEXPLORED: SCIENCE AND APPLICATIONS OF MEDICINAL PLANTS

JUNE 28 - JULY 2, 2022 | PUERTO RICO, USA

Saludos:

Human, animal and plant health, climate change, and food security remain among the greatest challenges for societies around the globe. Natural products and medicinally active plants can contribute significantly to the improvement of health. The 11th annual conference of the American Council for Medicinally Active Plants (ACMAP) invites you to join us to share scientific studies and lessons from lab and field to illustrate the latest advances in knowledge that can be used to impact and transform the lives of fellow human beings all over the world.

This year's meeting will focus on the scientific advances in several health and nutritional areas including the emerging research field of hemp and medicinal cannabis. Let this meeting be a forum where we come together to reflect on both basic and applied research and development activities that demonstrate how the application of science can and does make a difference to communities and society. It is a great pleasure in inviting you to present your research in Puerto Rico.

We look forward to a robust scientific discussion in Puerto Rico.

**Alok Arun (Chair, ACPMAP 2022)**

**Nirmal Joshee (Co-Chair, ACPMAP 2022)**



### HOSTS 2022



**JUAN NEGRON-BERRIOS, PH.D.**

HOST UNIVERSITY CHANCELLOR



**CARLOS OLIVARES, PH.D.**

HOST UNIVERSITY CHANCELLOR



**ALOK ARUN, PH.D.**

CHAIR



**NIRMAL JOSHEE, PH.D.**

CO-CHAIR

# 11<sup>th</sup> ANNUAL HYBRID CONFERENCE



## American Council for Medicinally Active Plants (ACMAP)

EXPLORING THE UNEXPLORED: SCIENCE AND APPLICATIONS OF MEDICINAL PLANTS

JUNE 28 - JULY 2, 2022 | PUERTO RICO, USA

¡Buenos días!

Dear Colleagues,

On behalf of the American Council for Medicinally Active Plants (ACMAP), I am honored and delighted to welcome you to the 11th Annual International Conference being held June 28 – July 2, 2022 at the Inter American University of Puerto Rico, Barranquitas, Puerto Rico. Last year's 10th Anniversary conference was held digitally due to the COVID-19 pandemic. We are overjoyed to be hosting this year's conference both in person and virtually.

The beauty of Puerto Rico's biological and cultural diversity has attracted many scientists who engage in multidisciplinary research, studying the botany and conservation of the unique species of Puerto Rico and the larger Caribbean, understanding their chemical structures and bioactivity, as well as the efficacy and safety of natural products for human health. The integrated efforts of researchers from diverse fields investigating the application of medicinally active plants will be required to translate basic science insights into public health benefit.

We are thrilled to officially open the conference with an outstanding Opening Plenary Lecture by Dr. David Julius, a 2021 Nobel Prize in Physiology and Medicine laureate. Dr. Julius's discoveries with capsaicin, the active component in red chili peppers, emphasize the tremendous potential of natural products research to develop breakthroughs that will prevent and treat many diseases. Dr. Julius's extraordinary work will inspire generations of students and young investigators involved in natural products research. This meeting has been made possible by the hard work of the organizing committee from the Inter American University of Puerto Rico and the ACMAP's Board of Directors. I am grateful for all the sponsors who provided support for the Conference.

Finally, I would like to thank all of you who are participating in the Conference for accepting our invitation and coming to Puerto Rico or joining us virtually to share our interest in medicinally active plants.

With best wishes for an outstanding meeting.

Anait S. Levenson, M.D., Ph.D.  
President, ACMAP

**ACMAP**



ANAIT S. LEVENSON, M.D., PH.D.  
LONG ISLAND UNIVERSITY, USA

# 11<sup>th</sup> ANNUAL HYBRID CONFERENCE



## American Council for Medicinally Active Plants (ACMAP)

EXPLORING THE UNEXPLORED: SCIENCE AND APPLICATIONS OF MEDICINAL PLANTS

JUNE 28 - JULY 2, 2022 | PUERTO RICO, USA

¡Bienvenidos!

On behalf of the American Council for Medicinally Active Plants (ACMAP), I am honored to welcome you to our 11th Annual Conference on the enchanting island of Puerto Rico. This year we are joined by presenters and attendees who are participating in-person and virtually from different parts of the world in Africa, Asia, Europe, North and South America, and the Caribbean.

ACMAP's goal is to discover new ways to link agriculture, biotechnology, ethnobotany, health, nutrition, and medicine and increase awareness of the need for plant-based medicinal research. To this end, this conference provides a forum to bring scientists, students, educators, and industry professionals to share their common interests. We hope you will exchange ideas and research findings that will stimulate the development of innovative means to produce medicinally active plants and identify new bioactive compounds with applications in human health.

I want to thank our Host and Chair, Dr. Alok Arun, and Co-Chair, Dr. Nirmal Joshee, and the local organization team at the Inter American University of Puerto Rico – Barranquitas for allowing us this opportunity to bring scientists together at this joint forum. Thank you for your attendance, and I invite you to enjoy a fruitful conference.

Sincerely,

Fabricio Medina-Bolivar, Ph.D.  
ACMAP Executive Director

**ACMAP**



FABRICIO MEDINA-BOLIVAR, PH.D.  
ARKANSAS STATE UNIVERSITY, USA

# 11<sup>th</sup> ANNUAL HYBRID CONFERENCE



## American Council for Medicinally Active Plants (ACMAP)

EXPLORING THE UNEXPLORED: SCIENCE AND APPLICATIONS OF MEDICINAL PLANTS

JUNE 28 - JULY 2, 2022 | PUERTO RICO, USA

### PLENARY SPEAKER

#### NOBEL LAUREATE



**DAVID JULIUS, PH.D.**

**UNIVERSITY OF CALIFORNIA  
SAN FRANCISCO, USA**

#### *From Peppers to Peppermints: Natural Products as Probes of the Pain Pathway*

Dr. David Julius used capsaicin, the active component in chili peppers, to identify the nerve sensors that allow the skin to respond to heat and pain.

Dr. David Julius was awarded the 2021 Nobel Prize in Physiology and Medicine jointly with Ardem Patapoutian for their discoveries of receptors for temperature and touch. Dr. Julius discovered molecular mechanisms of pain sensation and heat, including the characterization of the receptors that detect capsaicin, menthol, and temperature.

Dr. Julius earned his undergraduate degree from Massachusetts Institute of Technology and attained his doctorate from University of California, Berkeley. In 1997, his laboratory at the University of California, San Francisco cloned and characterized the transient receptor potential (TRPV1) channel, which is the receptor that detects capsaicin, the chemical in chili peppers that makes them "hot". Subsequently, Dr. Julius's laboratory contributed to the study of nociception by discovering other TRP channels that detect a range of temperatures and chemicals. These discoveries are vital to the development of therapeutics for chronic pain and other conditions.

<https://www.ibiology.org/neuroscience/sensory-receptors/>



# 11<sup>th</sup> ANNUAL HYBRID CONFERENCE

**American Council for Medicinally Active Plants (ACMAP)**

**EXPLORING THE UNEXPLORED: SCIENCE AND APPLICATIONS OF MEDICINAL PLANTS**

**JUNE 28 - JULY 2, 2022 | PUERTO RICO, USA**

## INVITED SPEAKERS



**ANAIT S. LEVENSON, M.D., PH.D.**  
LONG ISLAND UNIVERSITY, USA



**SUN-OK LEE, PH.D.**  
UNIVERSITY OF ARKANSAS, USA



**FABRICIO MEDINA-BOLIVAR, PH.D.**  
ARKANSAS STATE UNIVERSITY, USA



**JEFFREY ADELBERG, PH.D.**  
CLEMSON UNIVERSITY, USA



**JOSHUA KELLOGG, PH.D.**  
PENNSYLVANIA STATE UNIVERSITY, USA



**SRINIVASA RAO MENTREDDY, PH.D.**  
ALABAMA A&M UNIVERSITY, USA



**ANDREA I. DOSEFF, PH.D.**  
MICHIGAN STATE UNIVERSITY, USA



**JEREMY JOHNSON, PH.D.**  
UNIVERSITY OF ILLINOIS AT CHICAGO, USA



**YUANYUAN (ROSE) LI, PH.D.**  
UNIVERSITY OF MISSOURI, USA



**NIRMAL JOSHEE, PH.D.**  
FORT VALLEY STATE UNIVERSITY, USA



**OLIVER KAYSER, PH.D.**  
TU DORTMUND UNIVERSITY, GERMANY



**BAHAR ALIAKBARIAN, PH.D.**  
MICHIGAN STATE UNIVERSITY, USA



# INVITED SPEAKERS



**PRABODH SATYAL, PH.D.**

AROMATIC PLANT RESEARCH CENTER, USA



**NIHAL AHMAD, PH.D.**

UNIVERSITY OF WISCONSIN, USA



**SADANAND DHEKNEY, PH.D.**

UNIVERSITY OF MARYLAND  
EASTERN SHORE, USA



**NAMRITA LALL, PH.D.**

UNIVERSITY OF PRETORIA, SOUTH AFRICA



**MAHIPAL SHEKHAWAT, PH.D.**

GOVERNMENT INSTITUTE OF POSTGRADUATE  
STUDIES AND RESEARCH, INDIA



**VIRGINIE AIRES, PH.D.**

UNIVERSITY OF BURGUNDY, FRANCE



**SONIA PETER, PH.D.**

BIOCULTURAL EDUCATION AND  
RESEARCH PROGRAM, BARBADOS



**SYLVIA MITCHELL, PH.D.**

UNIVERSITY OF WEST INDIES, JAMAICA



**MARIA VILLARREAL ORTEGA, PH.D.**

UNIVERSIDAD AUTONOMA DEL  
ESTADO DE MORELOS, MEXICO



**DIANA ROOPCHAND, PH.D.**

RUTGERS UNIVERSITY, USA



**THOMAS TERRILL, PH.D.**

FORT VALLEY STATE UNIVERSITY, USA



**RODOLFO JULIANI, PH.D.**

RUTGERS UNIVERSITY, USA



**NIIRANJAN ARYAL, PH.D.**

VRX LABS, USA



**ARUNA WEERASOORIYA, PH.D.**

PRAIRIE VIEW A&M UNIVERSITY, USA



# **11<sup>th</sup> ANNUAL HYBRID CONFERENCE**

**American Council for Medicinally Active Plants (ACMAP)**

**EXPLORING THE UNEXPLORED: SCIENCE AND APPLICATIONS OF MEDICINAL PLANTS**

**JUNE 28 - JULY 2, 2022 | PUERTO RICO, USA**

## **TOPICS**



**Session I: Botanicals of Caribbean Diaspora**



**Session II: Bioactive Compounds of Plants and Their Roles in Human Health I (Cancer)**



**Session III: Chemistry of Medicinal Plants**



**Session IV: Ethnobotany, Bioprospecting and Production**



**Session V: In vitro Propagation and Conservation of Value-Added Plants**



**Session VI: Bioactive Compounds of Plants and Their Roles in Human Health II (Inflammatory Diseases)**



**Session VII: Hemp and Medical Cannabis**



**Session VIII: Plant and Algae Derived Cosmeceuticals and Nutraceuticals**





**11<sup>th</sup> ANNUAL HYBRID CONFERENCE**  
**American Council for Medicinally Active Plants (ACMAP)**  
**EXPLORING THE UNEXPLORED: SCIENCE AND APPLICATIONS OF MEDICINAL PLANTS**  
**JUNE 28 - JULY 2, 2022 | PUERTO RICO, USA**

## SCIENTIFIC PROGRAM

*Times are in Atlantic Standard Time (AST)*

Day 1 Inter American University of Puerto Rico (IAUPR) - Barranquitas Campus VENUE: MULTIUSOS Tuesday, June 28, 2022		
2:00 PM – 8:00 PM		COVID-19 Vaccination check and Registration
2:00 PM		Shuttle pick-up from Hyatt Place Hotel – Bayamon (outside the entrance door)
4:30 PM – 5:30 PM	Welcome	Master of Ceremony and Chair: Dr. Alok Arun Invocation: Rev. Arnaldo Cintron Miranda Co-Chair of ACMAP 2022: Dr. Nirmal Joshee Rector, Inter American University of Puerto Rico, Barranquitas Rector, Inter American University of Puerto Rico, Bayamon President, Inter American University of Puerto Rico Chairman, Board of Trustees, Inter American University of Puerto Rico Message of President ACMAP: Dr. Anait S. Levenson Message of Executive Director ACMAP: Dr. Fabricio Medina-Bolivar
5:30 PM – 6:15 PM	Plenary	Nobel Laureate Dr. David Julius (virtual) Morris Herzstein Chair in Molecular Biology and Medicine University of California San Francisco, USA  From peppers to peppermints: natural products as probes of the pain pathway  Q&A (moderated by Alok Arun)
6:15 PM – 7:30 PM	Opening ceremony	Cultural event and dinner  Vote of thanks: Alok Arun (Chair)  Group Photo
8:00 PM		Shuttle leave for Hyatt Hotel Bayamon





**11<sup>th</sup> ANNUAL HYBRID CONFERENCE**  
**American Council for Medicinally Active Plants (ACMAP)**  
**EXPLORING THE UNEXPLORED: SCIENCE AND APPLICATIONS OF MEDICINAL PLANTS**  
**JUNE 28 - JULY 2, 2022 | PUERTO RICO, USA**

## SCIENTIFIC PROGRAM

*Times are in Atlantic Standard Time (AST)*

Day 2 IAUPR – Bayamon Campus Wednesday, June 29, 2022			
Shuttle pick-up at 7:30 am (Hyatt Place Hotel – Bayamon)			
8:00 AM – 8:30 AM BREAKFAST			
	Chair  Co-chair	<b>Session I: Botanicals of Caribbean diaspora</b> <i>(Venue: Salon Usos Multiples/SUM)</i>  <b>S. Rao Mentreddy</b> Alabama A&M University, USA  <b>James Ackerman</b> University of Puerto Rico, Puerto Rico	<b>Session II: Bioactive compounds of plants and their roles in human health I (Cancer)</b> <i>(Venue: Theatre)</i>  <b>Andrea Doseff</b> Michigan State University, USA  <b>Claudia Ospina</b> Inter American University of Puerto Rico, Puerto Rico
8:30 AM – 9:10 AM	Keynote	<b>Sonia Peters</b> Bioscience Barbados Ltd., Barbados  Caribbean Plant Biodiversity – unlocking the potential of molecular diversity.	<b>Nihal Ahmad</b> University of Wisconsin, USA  Grape antioxidants in skin cancer management
9:10 AM – 9:40 AM	Invited I	<b>Sylvia Mitchell (virtual)</b> University of West Indies, Jamaica  Thirty-five years of tropical plant tissue culture R&D in Jamaica: the challenges and breakthroughs	<b>Anait S. Levenson</b> Long Island University, USA  The role of dietary stilbenes in prostate cancer interception: evidence from preclinical studies.
9:40 AM – 10:10 AM	Invited II	<b>Martha Giraldo</b> University of Puerto Rico, Puerto Rico  Role of roots and tuber crops in global household food security	<b>Yuanyuan (Rose) Li</b> University of Missouri, USA  Maternal epigenetic diets on transgenerational breast cancer prevention
10:10 AM – 10:40 AM	Invited III	<b>Alok Arun</b> Inter American University of Puerto Rico, Puerto Rico  Biology of underutilized crops of the Caribbean	<b>Virginie Aires (virtual)</b> University of Burgundy, France  Bioactive dietary compounds: promising therapeutic adjuvants through the targeting of tumor lipid metabolism
10:40AM – 11:00 AM	COFFEE BREAK		



**11<sup>th</sup> ANNUAL HYBRID CONFERENCE**  
**American Council for Medicinally Active Plants (ACMAP)**  
 EXPLORING THE UNEXPLORED: SCIENCE AND APPLICATIONS OF MEDICINAL PLANTS  
 JUNE 28 - JULY 2, 2022 | PUERTO RICO, USA

# SCIENTIFIC PROGRAM

*Times are in Atlantic Standard Time (AST)*

11:00 AM – 11:15 AM		Please move to Theatre	<b>Gisella Campanelli (virtual)</b> Long Island University, USA  Chemopreventive activity of Gnetin C in murine tumor models of prostate cancer
11:15 AM – 11:30 AM		Please move to Theatre	<b>Mirielle Nauman</b> University of Illinois at Chicago, USA  Garcinia mangostana (mangosteen) xanthones promote androgen receptor degradation
11:30 AM -12:30 PM	Workshop	Plant tissue culture of medicinally active plants (Jeffrey Adelberg, Sadanand Dhekney, Nirmal Joshee) (Venue: Salon Usos Multiples/SUM)	
12:20 PM- 1:30 PM	LUNCH		
	Chair	<b>Session III: Chemistry of medicinal Plants</b> <i>(Venue: Salon Usos Multiples/SUM)</i>  <b>Sun-Ok Lee</b> University of Arkansas, USA	<b>Session IV: Ethnobotany, bioprospecting and production</b> <i>(Venue: Theatre)</i>  <b>Jeffrey Adelberg</b> Clemson University, USA
	Co-chair	<b>Bruyanelis Ramos Aponte</b> Universidad Ana G. Méndez, Puerto Rico	<b>Angel Nunez Marrero</b> Inter American University of Puerto, Puerto Rico
1:30 PM -2:10 PM	Keynote	<b>Jeremy Johnson</b> University of Illinois at Chicago, USA  Translational development of phytochemicals and highly characterized plant extracts	<b>James E Simon (virtual)</b> Rutgers University, USA  Bioexploration of indigenous plants in sub-Sahara Africa
2:10 PM – 2:40 PM	Invited I	<b>Fabricio Medina-Bolivar</b> Arkansas State University, USA  Combined elicitation and targeted metabolomics approach for the discovery of medicinally active prenylated stilbenoids in hairy root cultures of peanut and pigeon pea	<b>S. Rao Mentreddy</b> Alabama A&M University, USA  Ethnobotany and production of three high-value medicinally active herbs, basil, turmeric, and mount mint in North Alabama
2:40 PM – 3:10 PM	Invited II	<b>Joshua J Kellogg</b> Pennsylvania State University, USA  Biochemometric molecular networking to discover new aryl hydrocarbon receptor ligands from the diet	<b>Prabodh Satyal (Virtual)</b> Aromatic Plant Research Centre, USA  Instrumental challenges with essential oil adulteration.



**11<sup>th</sup> ANNUAL HYBRID CONFERENCE**  
**American Council for Medicinally Active Plants (ACMAP)**  
**EXPLORING THE UNEXPLORED: SCIENCE AND APPLICATIONS OF MEDICINAL PLANTS**  
**JUNE 28 - JULY 2, 2022 | PUERTO RICO, USA**

## SCIENTIFIC PROGRAM

*Times are in Atlantic Standard Time (AST)*

3:10 PM – 3:40 PM	Invited III	<b>Rodolfo Juliani</b> Rutgers University, USA  Quality, chemistry, and biological activity of the West African spice, grains of paradise (Aframomum melegueta)	<b>Nirmal Joshee</b> Fort Valley State University, USA  Ethnobotanical knowledge paving way for medicinal compounds and value- added products in Paulownia tree
3:40 PM – 4:00 PM	COFFEE BREAK		
4:00 PM – 4:15 PM	Oral talk	<b>Joseph Horzempa</b> West Liberty University, USA  Dillapole, a compound extracted from fennel, exhibits potential as an antibacterial chemotherapy by dampening virulence factor expression	<b>B.K. Biswas</b> Fort Valley State University, USA  Stevia and Turmeric, consumption can help for anti-cancer benefits, know how to grow and consume
4:15 PM – 4:30 PM	Oral talk	<b>Marco Nuno De Canha</b> University of Pretoria, South Africa  Harnessing the potential of two southern African plants for the treatment of acne vulgaris	Please move to Salon Usos Multiples/SUM
4:30 PM – 6:00 PM	POSTER PRESENTATION		
Day 3 IAUPR – Bayamon Campus Thursday, June 30, 2022			
Shuttle pick-up at 7:30 am (Hyatt Place Hotel – Bayamon)			
8:00 AM – 8:30 AM	BREAKFAST		
	Chair  Co-chair	<b>Session V: In vitro propagation and conservation of value-added plants</b> <i>(Venue: Salon Usos Multiples/SUM)</i>  <b>Nirmal Joshee</b> Fort Valley State University, USA  <b>Dr. Elizabeth Padilla Crespo</b> Inter American University of Puerto Rico, Puerto Rico	<b>Session VI: Bioactive compounds of plants and their roles in human health II (Inflammatory diseases)</b> <i>(Venue: Theatre)</i>  <b>Anait S. Levenson</b> Long Island University, USA  <b>Dr. Lizbeth Romero Perez</b> Inter American University of Puerto Rico, Puerto Rico
8:30 AM – 9:10 AM	Keynote	<b>Sadanand Dhekney</b> University of Maryland, USA	<b>Andrea Doseff</b> Michigan State University, USA





**11<sup>th</sup> ANNUAL HYBRID CONFERENCE**  
**American Council for Medicinally Active Plants (ACMAP)**  
**EXPLORING THE UNEXPLORED: SCIENCE AND APPLICATIONS OF MEDICINAL PLANTS**  
**JUNE 28 - JULY 2, 2022 | PUERTO RICO, USA**

# SCIENTIFIC PROGRAM

*Times are in Atlantic Standard Time (AST)*

		Optimizing in vitro culture techniques for precision breeding of perennial fruits: grapevine as a model	Flavonoids regulate macrophage immune function controlling cancer and halting obesity
9:10 AM – 9:40 AM	Invited I	<b>Jeff Adelberg</b> Clemson University, USA  Nutrients and lighting factors in controlling medicinal plant culture	<b>Susanne Mertens-Talcott (virtual)</b> Texas A&M University, USA  Interactions of tannins with the intestinal microbiome and anti-inflammatory activities
9:40 AM – 10:10 AM	Invited II	<b>Mahipal Singh Shekhawat (virtual)</b> Puducherry, India  In vitro regeneration of medicinal plants and analysis of structural plasticity of cells and tissues towards changed environmental conditions: A way forward to develop climate-resilient plants	<b>Diana Roopchand</b> Rutgers University, USA  Grape polyphenols and cannabidiol: mechanisms for promoting metabolic resilience
10:10 AM – 10:40 AM	Invited III	<b>Maria Luisa Villarreal Ortega (virtual)</b> Universidad Autónoma del Estado de Morelos, Mexico  Conservation of Mexican medicinal species with sedative and anti-inflammatory properties	<b>Kevin Tveter (10:10 AM–10:25 AM)</b> Rutgers University, USA  Metabolic benefits of grape polyphenol consumption on bile acid signaling in mice with intestine- and liver-specific deletion of Fxr
			<b>Meenakshi Sudhakaran (10:25 AM–10:40 AM)</b> Michigan State University, USA  Flavone apigenin regulates innate immune response through RNA binding protein hnRNP A2
10:40 AM-11:00 AM	COFFEE BREAK		
11:00 AM-11:15 AM	Oral talk	<b>Ramana Gosukonda</b> Fort Valley State University, USA  Machine learning models in plant tissue culture: a case study of Brahmi	<b>Please move to Salon Usos Multiples/SUM</b>
11:15 AM – 11:30 AM	Oral talk	<b>Samantha Sherman (virtual)</b> Fort Valley State University, USA  Histological examination of in vitro hyperhydric cultures of <i>Scutellaria lateriflora</i> .	<b>Please move to Salon Usos Multiples/SUM</b>
11:30 AM -12:30 PM	Workshop	<b>Writing and submitting scientific papers to JMAP (Adolfina Koroch)</b> (Venue: Salon Usos Multiples/SUM)	





**11<sup>th</sup> ANNUAL HYBRID CONFERENCE**  
**American Council for Medicinally Active Plants (ACMAP)**  
**EXPLORING THE UNEXPLORED: SCIENCE AND APPLICATIONS OF MEDICINAL PLANTS**  
**JUNE 28 - JULY 2, 2022 | PUERTO RICO, USA**

# SCIENTIFIC PROGRAM

*Times are in Atlantic Standard Time (AST)*

12.30 PM – 1.30 PM	LUNCH		
1.30 PM – 6.30 PM	TOUR	Guided Old San Juan Tour (Contact registration desk for more information)	
Day 4 IAUPR Bayamon Campus Friday, July 1, 2022			
Shuttle pick-up at 7:30 am (Hyatt Place Hotel – Bayamon)			
8:00 AM – 8:30 AM	BREAKFAST		
	Chair	Session VII: Hemp and Medical Cannabis (Venue: Salon Usos Multiples/SUM)  Joshua Kellogg Fort Valley State University, USA	Session VIII: Plant and algae derived cosmeceuticals and nutraceuticals  Alok Arun Inter American University of Puerto Rico, Puerto Rico
8:30 AM – 9:10 AM	Keynote	Aruna Weerasooriya Prairie View A & M University, USA  Genetic complexity in Industrial hemp germplasm	Namrita Lall (virtual) University of Pretoria, South Africa  Reinforce urban skin by connecting with the origins
9:10 AM – 9:40 AM	Invited I	Oliver Kayser (virtual) TU Dortmund University, Germany  Endocannabinoid receptor pharmacology of rare cannabinoids	Sun-Ok Lee University of Arkansas, USA  Health-promoting effects of functional foods
9:40 AM – 10:10 AM	Invited II	Sadanand Dhekney University of Maryland Eastern Shore, USA  Screening industrial hemp cultivars for cannabinoid production in the Mid-Atlantic region	Bahar Aliakbarian (virtual) Michigan State University, USA  Upcycling food process byproducts for cosmeceutical applications: Case of cherry waste
10:10 AM – 10:40 AM	Invited III	Wesley Raup-Konsavage Penn State University, USA  Chasing the entourage effect in Cannabis	Thomas H. Terrill Fort Valley State University, USA  Nutraceutical and cosmeceutical properties of sericea lespedeza (Lespedeza cuneata)
10:40AM – 11:00 AM	COFFEE BREAK		
11:00 AM – 11:15 AM	Oral talk	Niranjan Aryal VRX Labs, USA	Ana-Marie Reid (virtual) University of Pretoria, South Africa





**11<sup>th</sup> ANNUAL HYBRID CONFERENCE**  
**American Council for Medicinally Active Plants (ACMAP)**  
**EXPLORING THE UNEXPLORED: SCIENCE AND APPLICATIONS OF MEDICINAL PLANTS**  
**JUNE 28 - JULY 2, 2022 | PUERTO RICO, USA**

## SCIENTIFIC PROGRAM

*Times are in Atlantic Standard Time (AST)*

		Cannabis testing labs; Quality control and standardization	Euclea natalensis A. DC., as an adjuvant, immunomodulator and hepatoprotectant
11:15 AM – 11:30 AM	Oral talk	<b>Ke Sui</b> Rutgers University, USA  Cannabidiol improves inflammation, glucose metabolism and bone loss in a murine model of female post-menopause in association with a bloom of Lactobacillus sp.	<b>Sepideh Mohammadhosseinpour (virtual)</b> Arkansas State University, USA  Prenylated stilbenoids from peanut increase the anticancer efficacy of paclitaxel in triple-negative breast cancer cells
11:30 AM – 12:10 PM	Keynote	Please move to Theatre	<b>A. Douglas Kinghorn (virtual)</b> The Ohio State University, USA  Discovery of anticancer agents of diverse natural origin
12:10 PM – 12:15 PM	Flash talk	Please move to Theatre	<b>Esther Mezhibovsky</b> Rutgers University, USA  Grape polyphenols may protect from diet induced obesity by preventing high-fat diet dampening of the hypothalamic-pituitary axis
12:15 PM – 12:30 PM	Flash talk	Please move to Theatre	<b>Claudia A. Ospina-Millán</b> Inter American University of Puerto Rico, Puerto Rico  Natural Products Research in Puerto Rico: Isolation of bioactive metabolites from tropical plants
12:30 PM – 1:30 PM	LUNCH		
1:30 PM – 4:30 PM	LEISURE		
4:30PM	Shuttle Pick for Restaurant from Hyatt Bayamon		
5:00 PM – 8:00 PM	CLOSING CEREMONY (GALA DINNER) AND AWARDS		
	Day 5 Saturday, July 2, 2022 Field Excursions and Tours (Group basis) Contact Registration/Information desk for details		





# SESSION I: BOTANICALS OF CARIBBEAN DIASPORA

## KEYNOTE

### 01. Caribbean Plant Biodiversity - unlocking the potential of molecular diversity

Sonia Peter<sup>1</sup>, John Headley<sup>2</sup>, Neelam Buxani<sup>3</sup>. <sup>1</sup>Biocultural Education and Research Programme, 59 Meadowvale, St. James, Barbados, BB23022;

<sup>2</sup>National Hydrology Research Centre, Environment Canada, 11 Innovation Boulevard Saskatoon, SK S7N 3H5, Canada; <sup>3</sup>Clinical Development & Medical Affairs (CDMA), Roche Diagnostics Solutions, 4300 Hacienda Drive, Pleasanton, CA 94588 USA. E-mail: biosciencebdos@gmail.com

Plant biodiversity is significant in the Caribbean with the estimated 11,000 species having a high endemism of 72%. Of the estimated 1520 genera, 192 are considered endemic (Critical Ecosystem Partnership Fund). Endemic plant genera are concentrated in the Greater Antilles with greatest diversity recorded for the larger and more heterogeneous islands of Hispaniola and Cuba. The five plant families with the highest number of endemic genera are Asteraceae (39), Rubiaceae (27), Orchidaceae (12), Euphorbiaceae (11) and Fabaceae (10). Protection of this resource is vital as it has been a key source of survival for Caribbean populations providing food, shelter, textiles, and medicine. Unique genera possess phytochemical scaffolds that lead to the biosynthesis of novel bioactive agents in many natural product classes including the heterocyclic polyphenolics and flavonoids. Chemotaxonomic features of polyphenolic complexity will also be influenced by chemotypes and biogeographical features including soil composition, altitude, and weather variability. Two families of global significance for bioactivity are the Lamiaceae and Asteraceae. Though the Lamiaceae is not representative of diverse genera in the Caribbean, an examination of chemotaxonomic features in natural products from island Lamiaceae species reflects the molecular diversity that must be present within the Caribbean islands' endemic genera. An assessment of Lamiaceae species extracts via HPLC MS MS demonstrated the tendency for biosynthesis of flavonoids, and flavonoid glycosides, with methoxylation on rings A, B and C of the flavonoid skeleton. The application of computer modelling to further unlock the bioactivity potential of these natural agents demonstrated the potential for molecular intervention against the SARS COV 2 corona virus, as a biological target. Molecular modelling has become a fast-track tool to clinical trials of bioactive compounds. The molecular diversity within the Caribbean archipelago is deserving of concerted focus as we search for novel bioactive agents.

# SESSION I: BOTANICALS OF CARIBBEAN DIASPORA

## INVITED TALK

### O2. Thirty-five years of tropical plant tissue culture R&D in Jamaica: the challenges and breakthroughs

Sylvia Adjoa Mitchell. Medicinal Plant Research Group, The Biotechnology Centre, The University of the West Indies, Mona Campus, Jamaica.

E-mail: [sylvia.mitchell@uwimona.edu.jm](mailto:sylvia.mitchell@uwimona.edu.jm)

In 1985, the International Institute of Tropical Agriculture (IITA) initiated a joint GOJ/UWI/IITA Root Crop Project. This project introduced minisett yam technology to Jamaica, enabled tissue culture capabilities at Life Sciences, UWI and built a tissue culture facility at the Scientific Research Council. In 1989, another tissue culture facility was initiated at The Biotechnology Centre, UWI, courtesy of a grant from the European Communities. The Centre has maintained training and research facilitating postgraduates pursuing their MPhil or PhD in Biotechnology. The Medicinal Plant Research Group (MPRG) was initiated at the Centre in 1999. A five-prong approach has been refined by the MPRG to include 1) bioactivity (antifungal, antibacterial) and biochemical (chlorophyll level, antioxidant activity) studies of various crops (bottlebrush, breadfruit, jackfruit) which has resulted in several refereed papers and commercial business development; 2) ethnobotany studies and monographs; 3) tissue culture studies adding: bamboo (*Bambusa vulgaris*), bottle brush (*Callistemon viminalis*), cassava (*Manihot esculenta*), chainy root (*Smilax balbisiana*), pepper elder (*Piper amalago*), sweet potato (*Ipomea batatas*), and yams (*Dioscorea* spp); 4) business support including taxonomic review of Jamaican root tonic plants, TRAMIL Jamaican study, and data gathering support to several country-wide projects, 5) International links – Caribbean Access and Benefit Sharing for ratification of the Nagoya Protocol, IUCN report on Caribbean Bioprospecting, the SRAD research cluster; Edinburgh projects on reformatory justice and SOAD with links to Africa. There has been a conscious effort to be involved in and to tailor research to serve rural and industrial needs. There has also been a conscious effort to mix short-term research that has immediate application (eg. development of low-cost tissue culture kits, clean planting material) with longer-term R&D that may take years to apply but for which the potential is much greater (e.g., biofarming, somatic embryogenesis of elite trees, product development, commercial tissue culture). The challenges and opportunities arising from these activities will be discussed.

## 11<sup>th</sup> ANNUAL HYBRID CONFERENCE

**American Council for Medicinally Active Plants (ACMAP)**

EXPLORING THE UNEXPLORED: SCIENCE AND APPLICATIONS OF MEDICINAL PLANTS

JUNE 28 - JULY 2, 2022 | PUERTO RICO, USA



## **SESSION I:**

# **BOTANICALS OF CARIBBEAN DIASPORA**

### **INVITED TALK**

#### **O3. Role of roots and tuber crops in global household food security**

Martha. C. Giraldo. Agroenvironmental Sciences Department, University of Puerto Rico Mayagüez Campus.

Email: [martha.giraldo@upr.edu](mailto:martha.giraldo@upr.edu)

Root crops are part of the regular diets in Puerto Rico, the Caribbean, and the Southern US. The sweet potato, yam, and tanier occupy an essential place as a primary food source. As part of global climate change and hurricane seasons is imperative for the South US, Caribbean, and Puerto Rico to strengthen their food security sources, and one of our main assets relies on our root crops. These root crops have not only the resilience for production but the nutritional facts required. Even though these food crops are mostly seen as minor or specialty crops, they have the potential to feed and be part of the solution to reinforce food security for the Caribbean and worldwide. Concerning yam, tanier, and sweet potato crops, as of 2019, it was reported that their value at the farm level is close to \$6 million, with yam being the most economically important, followed by tanier and sweet potato. The contribution of these to total consumption fluctuated from 3.6% in sweet potato, 7% in tanier, and 20% in yam. We have a significant margin for growth in these crops, which are so important for local nourishing. These crops can significantly improve nutrition and food security. As essential cash crops, they present several mutual challenges. They are propagated clonally by vegetative material, allowing yield-reducing pathogens and pests to disperse and build up over time. This talk aims to address important research projects that are currently being developed to achieve this goal. Achieving greater food security is important for Puerto Rico and the world.

**11<sup>th</sup> ANNUAL HYBRID CONFERENCE**

**American Council for Medicinally Active Plants (ACMAP)**

**EXPLORING THE UNEXPLORED: SCIENCE AND APPLICATIONS OF MEDICINAL PLANTS**

**JUNE 28 - JULY 2, 2022 | PUERTO RICO, USA**



# SESSION I: BOTANICALS OF CARIBBEAN DIASPORA

## INVITED TALK

### O4. Development of scientific tools for underutilized crops of Caribbean

Alok Arun<sup>1\*</sup>, Celine Noirot<sup>2</sup>, Rosalinda Aybar Batista<sup>1</sup>, Jacobo Morales Mendez<sup>1</sup>, Juan Colon Morales<sup>1</sup>, Natalia Berríos Rivera<sup>1</sup>, & Juan Negrón Berríos<sup>1</sup>. <sup>1</sup>Institute of Sustainable Biotechnology, Department of Science and Technology, Inter American University of Puerto Rico, Barranquitas, PO Box 517, Puerto Rico, USA . <sup>2</sup>224 Chemin de Borde Rouge - Auzeville CS 52 627 - 31326 Castanet Tolosan cedex, France. Email: alok\_arun@br.inter.edu

The Peruvian carrot (*Arracacia xanthorrhiza*) commonly referred, as apio is a crop that is a secondary food item for over 100 million people, mainly in South America and Puerto Rico. While the storage roots are the main product, the rootstock and leaves are used as animal feed and the aerial stems are used as propagules. The crop has low input requirements and can be grown in a variety of frost-free tropical highland environments. However, in the absence of preventive management practices and proper handling, infections caused by pathogens severely damage the storage roots. The lack of certainty about the agents and the process of infection is a major gap to understanding how to prevent infection at its very onset. Despite the economical and agricultural importance of apio, any biotechnological strategy aimed at improving the yield or protecting the crop from pest damage is non-existent, partly due to lack of any genomic tools and techniques. For non-model plants like apio that has very limited genomic information available, developing genomic and transcriptome databases may help improve its cultivation and shelf life. In our study, we characterized the global gene expression profile of *A. xanthorrhiza* tissues (tuber, leaf, flower and cormel) using Illumina high-throughput RNA sequencing platform. The study generated 74,533 contigs corresponding to approximately 72 million paired-end reads. Functional annotation of the dataset identified 12 of the 15 homologous enzymes involved in flavonoid biosynthesis pathway in plants and six gene families involved in starch biosynthesis. Further, we sequenced the nuclear genome of *Arracacia xanthorrhiza* using PacBioscience platform, which generated 117,654, reads with mean length of 61,758 bps. Currently, we are assembling and annotating the genome. Finally, we studied the tissue structure of the tuber using Scanning Electron Microscope to understand the structure of starch molecules. The present study is the maiden report on transcriptome and genome datasets in *Arracacia* species and dataset will provide a resource for further research on improving the quality and storage life of the crop.

Acknowledgment: USDA HSI STEM grant no. 2021-77040-34872 and UIPR Fondo Semilla grant awarded to Alok Arun supported this research.



## SESSION II:

# BIOACTIVE COMPOUNDS OF PLANTS AND THEIR ROLES IN HUMAN HEALTH I (CANCER)

### KEYNOTE

#### O5. Grape antioxidants in skin cancer management

Chandra K. Singh<sup>1</sup> and Nihal Ahmad<sup>2</sup>. <sup>1</sup>Department of Dermatology, University of Wisconsin, and <sup>2</sup>William S. Middleton VA Hospital, Madison, WI.  
E-mail: nahmad@dermatology.wisc.edu

Non-melanoma skin cancers (NMSCs) are the most frequently diagnosed cancers in the USA, with an estimated 5.4 million cases diagnosed annually in 3.3 million people in the USA. Therefore novel approaches are needed for the management of NMSCs. Dietary agents have gained considerable attention for protection against skin cancer. Red grapes contain a natural amalgamation of agents with antioxidant and anti-inflammatory properties. We have found that GP has significant chemopreventive effects against ultraviolet B (UVB) radiation mediated cutaneous responses, including photocarcinogenesis in male and female SKH1 hairless mice. The observed effects of GP were associated with modulations in multiple cellular processes (DNA repair, oxidative stress, protein ubiquitination and acute phase response signaling), in female mice. Similarly, we found a marked reduction in malignant conversion of premalignant lesions in GP-treated male mice. Additionally, we observed significant decrease in mast cell infiltration and serum IgE and cytokine CCL1 levels. To further understand the mechanisms of the biological response of GP, we employed cancer-pathway PCR array to profile 84 key genes related to transformation and tumorigenesis, in tissues obtained from mice. GP feeding modulated several genes related to cellular senescence, DNA repair, hypoxia and telomere maintenance. We observed significant modulation in 6 genes involved in hypoxia signaling (Hmox1, and Ldha), DNA repair (Ercc3), and telomeres and telomerase (Terf2ip, Tinf and Tnks2). Ingenuity Pathway Analysis (IPA) suggested association with increased cellular homeostasis and apoptosis, as well as decreased cell migration, viability and proliferation in tumors, which all support the chemoprotective response of GP. Additionally, IPA suggested probable inhibition of H2O2, PI3K, STAT3, IGF1, IL1A, IL1B, RELA, and TP53 activation, as the upstream regulators of GP-altered genes. A network pathway of interacting genes further suggested links to key players in skin cancer (NF-κB, PI3K, P38 MAPK, ERK1/2, AMPK, CD3, and histones H3 and H4) supporting anti-tumor properties of dietary grape. Overall, our studies suggest that grape consumption has chemopreventive effects against skin cancer.

## **SESSION II: BIOACTIVE COMPOUNDS OF PLANTS AND THEIR ROLES IN HUMAN HEALTH I (CANCER)**

### **INVITED TALK**

#### **O6. The role of dietary stilbenes in prostate cancer interception: evidence from preclinical studies**

Anait S Levenson. Department of Biomedical Sciences, College of Veterinary Medicine, Long Island University, Brookville, NY 11548, USA.

E-mail: [anait.levenson@liu.edu](mailto:anait.levenson@liu.edu)

Developing novel interceptive strategies for prostate cancer management is a supreme goal in clinical oncology. Polyphenols found in food and plants, particularly stilbenes, have gained increased importance as potential preventive and therapeutic anticancer agents due to their potent biological properties including anti-inflammatory, anti-oxidative, anti-proliferative, and pro-apoptotic effects. Studies have shown that stilbenes can affect multiple signaling pathways involved in inflammation, cell proliferation and survival, EMT, invasion, angiogenesis, and metastasis. In fact, stilbenes are shown to modulate epigenetic and transcription factors, growth factors, enzymes, and non-coding RNAs. In our studies, we demonstrate metastasis associated protein 1 (MTA1)-mediated anti-inflammatory and anticancer effects of resveratrol, pterostilbene and Gnetin C in prostate cancer in vitro and in vivo. Importantly, we conducted various prostate cancer pre-clinical studies with stilbenes using both transgenic mouse models and xenografts and demonstrated potent MTA1-targeted interceptive outcome. I will discuss the importance of mechanistic preclinical studies with natural products for future successful human clinical trials towards protecting select responsive group of patients.



## **SESSION II:**

# **BIOACTIVE COMPOUNDS OF PLANTS AND THEIR ROLES IN HUMAN HEALTH I (CANCER)**

### **INVITED TALK**

#### **O7. Maternal epigenetic diets on transgenerational breast cancer prevention**

Yuanyuan (Rose) Li\*. Department of Obstetrics, Gynecology & Women's Health, Department of Surgery, University of Missouri \*Address correspondence to Yuanyuan (Rose) Li, One Hospital Drive, Medical Sciences Building, M659, Columbia, MO, 65212, Tel: +1-573-884-8975, E-mail: ylgrk@missouri.edu.

Breast cancer is the most common type of cancer and the second leading cause of death among women in the United States. Studies have shown breast cancer has strong developmental origins. Maternal nutrition has a significant impact on epigenetic reprogramming processes during early development, which may alter the offspring's health outcome later in adult life. In the current studies, we seek direct evidence that development of breast cancer may originate from a fetal environment and maternal nutrition composition may influence the disease outcome through epigenetic mechanisms. Our results showed that the prevention outcome for maternal soybean genistein (GE) was explicitly dependent on exposure timing, and temporal epigenetic changes controlled important gene expression that are central to the efficacy of maternal GE on breast cancer prevention in later life. This study provides important insights into an appropriate maternal administration of soybean-based botanical diets or bioactive compounds that can maximize their beneficial effects leading to improved health outcome such as breast cancer prevention later in offspring life.

## SESSION II:

# BIOACTIVE COMPOUNDS OF PLANTS AND THEIR ROLES IN HUMAN HEALTH I (CANCER)

### INVITED TALK

#### O8. Bioactive dietary compounds : promising therapeutic adjuvants through the targeting of tumor lipid metabolism

Virginie Aires <sup>1, 2</sup>. <sup>1</sup>University of Burgundy, Dijon, F-21000, France ; <sup>2</sup>Inserm Research Center, URM 1231 "Lipids, Nutrition, Cancer (LNC)" – "Cancer and Adaptive Immune Response (CADIR)" team, "Bioactive Molecules and Health (BMH)" research group. E-mail: virginie.aires02@u-bourgogne.fr

Lipid metabolic reprogramming has now been largely recognized as a hallmark of cancer progression and metastasis. Among the rewired lipid metabolic programmes, that of phosphatidylcholine (PC, a major phospholipid of biological membranes) appears to be a common trait in many cancer types. We recently showed that altered PC metabolism supports colorectal cancer (CRC) cells resistance to chemotherapies such as 5-Fluorouracil (5-FU). Indeed, altered PC metabolism in CRC models was associated to an increase in the expression and activity of an enzyme, lysophosphatidylcholine acyltransferase 2 (LPCAT2), which supports intracellular PC synthesis and the production of intracellular organelles called lipid droplets (LD). In collaboration with the Georges François Leclerc Anticancer Centre in Dijon, immunohistochemical analyses of the expression of LPCAT2 on liver metastases sections from 54 CRC patients who received neoadjuvant chemotherapy, suggests a correlation between LPCAT2 expression, response to treatment and relapse-free survival. The latter could partly be explained by a modulation of the tumor microenvironment, characterized by decreased immunogenic death and a decreased CD8<sup>+</sup> T lymphocytes infiltration at the tumor site, in vivo in animals and in CRC patients with liver metastases. In view of the importance of PC metabolism in chemoresistance mechanisms, the use of compounds being able to modulate PC levels, may be a relevant strategy to re-sensitize tumor cells to treatments. Among these compounds, naturally occurring polyphenols may be good candidates. In this presentation, we will focus on the well-known polyphenol resveratrol, and show in various CRC cellular models that its chemosensitizing effects are partly due to its ability to modulate PC levels through the inhibition of key enzymes involved in both PC de novo and remodelling pathways.

This work was supported by grants from the Conseil Régional de Bourgogne-Franche-Comté, the FEDER, the Bureau Interprofessionnel des Vins de Bourgogne and the Laboratoire d'Excellence LipSTIC.



## SESSION II: BIOACTIVE COMPOUNDS OF PLANTS AND THEIR ROLES IN HUMAN HEALTH I (CANCER)

### CONTRIBUTED ORAL TALK

#### **09. Chemopreventive activity of Gnetin C in murine tumor models of prostate cancer**

Gisella Campanelli<sup>1</sup>, Prashanth Parupathi<sup>1</sup>, Qing Cai<sup>1</sup>, Avinash Kumar<sup>1</sup>, Anait S Levenson<sup>2</sup>. <sup>1</sup>Arnold & Marie Schwartz College of Pharmacy and Health Sciences, Long Island University, Brooklyn, NY 11201, USA; <sup>2</sup>School of Veterinary Medicine, Long Island University, Brookville, NY 11548, USA. E-mail: gisella.campanelli@my.liu.edu; anait.levenson@liu.edu

Prostate cancer (PCa) is the most common of all cancers in men. Epidemiological data suggest that diet has substantial impact on the development and progression of PCa. Among dietary polyphenols, stilbenes have been studied for their potential as chemopreventive and therapeutic agents in PCa. Gnetin C, a resveratrol-dimer found abundantly in melinjo plant (*Gnetum gnemon*), possesses more potent biological properties compared to resveratrol and pterostilbene (Pter) due, in part, to its improved pharmacokinetics. In our previous studies, we have extensively reported on the metastasis-associated protein 1 (MTA1)- and androgen receptor (AR) -targeted anti-inflammatory and anticancer properties of stilbenes. Here, for the first time, we examine the efficacy of Gnetin C in clinically adequate transgenic mouse models representing early and advanced stages of PCa progression. For this study, we generated prostate-specific MTA1-knocked in transgenes on the background of Pten heterozygosity (R26MTA1; Pten<sup>+/f</sup>; Pb-Cre<sup>+</sup>) and Pten knockout (R26MTA1; Pten<sup>f/f</sup>; Pb-Cre<sup>+</sup>). R26MTA1; Pten<sup>+/f</sup>; Pb-Cre<sup>+</sup> mice were used for chemopreventive strategy (diet supplementation with Gnetin C) while R26MTA1; Pten<sup>f/f</sup>; Pb-Cre<sup>+</sup> mice with more aggressive PCa were injected with Gnetin C as therapeutic strategy. Our results show that mice treated with Gnetin C (7 mg/kg bw, daily i.p.) exhibited more favorable histopathology with decreased severity and number of PIN foci (41.23 % reduction) accompanied by reduced proliferation (Ki67, 57.43% reduction), angiogenesis (CD31, 86.91% reduction), AR (31.57 % reduction), and MTA1 (27.94 % reduction). Ongoing chemopreventive studies with four different diets (Control-vehicle group; Pter (70 mg/kg diet); Gnetin C high dose (70 mg/kg diet); and Gnetin C low dose (35 mg/kg diet)) and are in progress. Taken together, these data provide an evidence for efficacy of Gnetin C in blocking PCa progression and implicate the potential of Gnetin C in PCa chemoprevention and therapy.

## SESSION II: BIOACTIVE COMPOUNDS OF PLANTS AND THEIR ROLES IN HUMAN HEALTH I (CANCER)

CONTRIBUTED ORAL TALK

### **O10. *Garcinia mangostana* (mangosteen) xanthones promote androgen receptor degradation**

Mirielle Nauman<sup>1,2</sup>, Jonghoon Won<sup>2</sup>, and Jeremy Johnson<sup>1,2</sup>. <sup>1</sup>Department of Pharmaceutical Sciences, University of Illinois at Chicago College of Pharmacy, Chicago, IL 60612, <sup>2</sup>Department of Pharmacy Practice, University of Illinois at Chicago College of Pharmacy, Chicago, IL 60612. E-mail addresses: mnauma6@uic.edu, jjjohn@uic.edu

The purple mangosteen fruit (*Garcinia mangostana*) contains a unique class of chemical compounds called the xanthones. Our lab has evaluated these xanthones for their potential activity in breast, colon, and prostate cancer cells. Preliminary data has shown that isoprenylated xanthones exhibit cytotoxic activity in multiple cell lines, specifically  $\alpha$ -mangostin and gartanin, two of the most abundant mangosteen xanthones. In prostate cancer cells, both compounds induce apoptosis and induce androgen receptor (AR) degradation. Mechanistic data reveals that there is a simultaneous decrease in AR protein expression and an increase in BiP, a cell stress chaperone, protein expression in prostate cancer cells treated with  $\alpha$ -mangostin. Ubiquitination of the androgen receptor via the proteasome has also been observed through western blots and immunoprecipitation experiments. Interestingly,  $\alpha$ -mangostin also promotes degradation of mutated AR, containing either point mutations or splice variants, both of which are mechanisms of drug resistance for prostate cancer cells and present significant clinical problems. In vivo data has confirmed that  $\alpha$ -mangostin is effective at reducing tumor size in mice xenografted with prostate cancer cells, even more effective than Bicalutamide, and that there may be in vivo modulation of the androgen receptor. We hypothesize that  $\alpha$ -mangostin promotes degradation of both wild type and mutated AR by promoting an interaction between AR and BiP proteins. This data highlights a class of compounds that targets AR in drug resistant prostate cancer cases through a novel and unique strategy.



## SESSION III: CHEMISTRY OF MEDICINAL PLANTS

### **O11. Translational development of phytochemicals and highly characterized plant extracts**

Jeremy J. Johnson. Department of Pharmacy Practice, University of Illinois Chicago, Chicago, IL 60612. E-mail: [jjjohn@uic.edu](mailto:jjjohn@uic.edu)

Dose selection and the pharmacokinetic properties of phytochemicals is critical in the successful translation of promising study agents. The traditional approach with plant based medicine has been to isolate the “active ingredient” from a plant and utilize a high dose to achieve the greatest activity. For certain situations this may be appropriate, however, in other circumstances this could actually hinder the development of a study agent. More recently, it is becoming increasingly clear that the whole plant matrix likely contributes to biological activity. In addition, evidence is suggesting that higher doses may not be optimal for improving the pharmacokinetic and pharmacodynamic endpoints of phytochemicals. Herein we will describe important considerations when deciding if a pure phytochemical or a complex plant matrix is the ideal choice as study agents as they progress to early phase clinical trials. From our own research as well others we will describe pharmacokinetic examples of several phytochemicals from different plants including rosemary (*Salvia rosmarinus*), green tea (*Camellia sinensis*), and mangosteen (*Garcinia mangostana*). Discussion will focus on dose selection and how raw material may impact the pharmacokinetic and pharmacodynamic endpoints in a pre-clinical and clinical research.

## **SESSION III: CHEMISTRY OF MEDICINAL PLANTS**

### **INVITED TALK**

#### **O12. Combined elicitation and targeted metabolomics approach for the discovery of medicinally active prenylated stilbenoids in hairy root cultures of peanut and pigeon pea**

Fabricio Medina-Bolivar. Department of Biological Sciences and Arkansas Biosciences Institute, Arkansas State University, Jonesboro, AR 72401, USA.  
E-mail: fmedinabolivar@astate.edu

Prenylated stilbenoids are a class of phytoalexins produced in certain plants such as peanut and pigeon pea. Studies to elucidate the biological activities of these compounds have been limited because they are not commercially available and are produced at very low levels in plants making their isolation challenging. To address these issues, we established hairy root cultures of peanut and pigeon pea and co-elicited them with methyl jasmonate, cyclodextrin, hydrogen peroxide and magnesium chloride to induce the biosynthesis of prenylated stilbenoids. In peanut, several prenylated stilbenoids accumulated in the culture medium. These included arachidin-1, arachidin-2, arachidin-3, arachidin-5, and arachidin-6. In pigeon pea, the main prenylated stilbenoid was cajaninstilbene acid. Interestingly, the stilbenoids identified in peanut represent analogs of resveratrol. Whereas the stilbenoids in pigeon pea represent analogs of pinosylvin. In both species, more than 95% of the total stilbenoids identified were found in the culture medium. Targeted tandem mass spectrometry analyses revealed that several other stilbenoids were also induced and secreted into the culture medium. To evaluate their biological activities, the prenylated stilbenoids from peanut were purified using column chromatography and semi-preparative high-performance liquid chromatography. The purified prenylated stilbenoids showed antioxidant, anti-inflammatory and anticancer activities in vitro. The hairy root cultures of peanut and pigeon pea demonstrated to be a reliable and sustainable platform for producing different types of prenylated stilbenoids with potential applications in human health.



## SESSION III: CHEMISTRY OF MEDICINAL PLANTS

### INVITED TALK

#### **O13. Biochemometric Molecular Networking to Discover New Aryl Hydrocarbon Receptor Ligands from the Diet**

Joshua J. Kellogg<sup>1,2</sup>. <sup>1</sup>Department of Veterinary and Biomedical Sciences and <sup>2</sup>Huck Institute of the Life Sciences, Pennsylvania State University, University Park, PA. E-mail: [jjk6146@psu.edu](mailto:jjk6146@psu.edu)

The aryl hydrocarbon receptor (AHR) has emerged as a critical ligand-activated transcription factor that promotes barrier tissue integrity, cellular differentiation programming, and immune system homeostasis, especially in the gastrointestinal tract, where the presence of AHR ligands enhances resistance to toxic insults. While botanical foods are known to exert AHR effects, the understanding of which specific bioactive molecules responsible for AHR activity is generally wanting. We developed a library of known AHR ligands to use as a reference database to screen for potential ligands from common foods (mushrooms, peppers, carrots). Using a molecular networking approach, combined with biochemometric modeling, we uncovered several compounds that possessed AHR-activity in an in vitro model. The network allowed for preliminary structural information about the dietary ligands, and demonstrated enhanced detection capabilities compared to previous studies that utilized conventional bioassay guided fractionation discovery. These results suggest that there are a variety of phytochemicals present in the diet that are capable of modulating AHR activity and potentially affecting downstream immune function in the gut.

## SESSION III: CHEMISTRY OF MEDICINAL PLANTS

### INVITED TALK

#### **O14. Quality, chemistry, and biological activity of the West African spice, grains of paradise (*Aframomum melegueta*)**

H. Rodolfo Juliani. Department of Plant Biology. School of Environmental and Biological Sciences. Rutgers University. 59 Dudley Rd. New Brunswick, 08901, New Jersey.

Herbs and spices have seen a significant increase in their use during the 21st century. Consumers have been searching for new ways to spice up their lives, looking for new experiences, products, flavors, and spices. This trend has increased during the COVID19 pandemic, as people needed to hunker down at home and then cook food and introduce variety, color, and new flavors to foods through spicing. Grains of paradise (*Aframomum melegueta*), a member of the ginger family (Zingiberaceae), originated from West Africa, specifically, the countries of the Guinea forest, Ghana, and Liberia. Grains of paradise is a seed spice, like cardamom. Grains of paradise contains essential oils and non-volatile components, both responsible for the aroma and flavor. This presentation aims to highlight the recent advances in the quality, chemistry, and biological activities of the West African spice. In terms of quality, developing standards can increase its commercial interest. As part of my research at Rutgers University, I contributed in collaboration with the US and African colleagues to define those standards of quality. The presentation will highlight recent activities on standard developments, including macroscopic, microscopic characterizations, chemical and proximate analysis. The presentation will focus on our recent work on the chemical diversity of non-volatile polyphenols of grains of paradise and new biological activities.



## SESSION III: CHEMISTRY OF MEDICINAL PLANTS

### CONTRIBUTED ORAL TALK

#### **O15. Dillapiole, a compound extracted from fennel, exhibits potential as an antibacterial chemotherapy by dampening virulence factor expression**

Elliot Collins<sup>1</sup>, Deanna M. Schmitt<sup>1</sup>, Francisco Leon<sup>2</sup>, Kristen Sikorsky<sup>1</sup>, Anthony Sako<sup>1</sup>, James Denvir<sup>3</sup>, Donald A. Primerano<sup>3</sup>, and Joseph Horzempa<sup>1</sup>.

<sup>1</sup>Department of Biomedical Sciences, West Liberty University, West Liberty, WV.

<sup>2</sup>Drug Discovery & Biomedical Sciences, College of Pharmacy, University of South Carolina, Columbia, SC. <sup>3</sup>Department of Biomedical Sciences, Marshall University, Huntington, WV E-mail: joseph.horzempa@westliberty.edu

*Francisella tularensis* has been classified as a Category A Bioterror Agent by the CDC. The intentional release of an antibiotic-resistant strain of *F. tularensis* could be devastating. Consequently, there is a need for novel treatments for *F. tularensis* infections. We previously tested a library of natural extracts to identify compounds that inhibit replication of a fluorescent *F. tularensis* strain during infection of THP-1 monocytes, but not outside the context of infection. One of the most promising extracts identified here was from Fennel (*Foeniculum vulgare*). Using bioassay guided fractionation, the Fennel extract was separated until a pure compound (dillapiole) was isolated and identified (via NMR and MS). Bona fide dillapiole (Sigma Aldrich) exhibited a similar effect to the purified compound, validating our findings. We first hypothesized that dillapiole may be augmenting host immunity. However, a multiplex analysis of RAW 264.7 macrophages suggested that only TNF- $\alpha$  was modulated by dillapiole. This finding was confirmed by ELISA. Moreover, RNA-seq data generated from THP-1 cells treated with dillapiole vs. vehicle were analyzed by Ingenuity Pathway Analysis. This analysis did not reveal an obvious host response that would explain the reduction in *F. tularensis* CFU upon treatment with dillapiole. Therefore, we hypothesized that dillapiole may be dampening virulence gene expression of *F. tularensis*. RNA-seq revealed that bacteria treated with dillapiole exhibited a reduction in expression of genes encoded in the *Francisella* Pathogenicity Island (FPI). Western blotting confirmed that dillapiole led to reduced production of IgA and IgC (two FPI proteins), but not LpnA (encoded elsewhere). This study shows that dillapiole is a promising novel type of antimicrobial that dampens virulence factor expression. Therefore, this compound does not exert the degree of selective pressure compared to traditional antibiotics which will likely minimize the development of resistance to dillapiole.

**11<sup>th</sup> ANNUAL HYBRID CONFERENCE**

**American Council for Medicinally Active Plants (ACMAP)**

**EXPLORING THE UNEXPLORED: SCIENCE AND APPLICATIONS OF MEDICINAL PLANTS**

**JUNE 28 - JULY 2, 2022 | PUERTO RICO, USA**



## SESSION III: CHEMISTRY OF MEDICINAL PLANTS

### CONTRIBUTED ORAL TALK

#### O16. Harnessing the potential of two southern African plants for the treatment of acne vulgaris

Marco Nuno De Canha<sup>1</sup> and Namrita Lall<sup>1,2,3</sup>. <sup>1</sup>Department of Plant and Soil Sciences, University of Pretoria, South Africa, Pretoria, 0002, <sup>2</sup>School of Natural Resources, University of Missouri, Columbia, MO 65211, USA and <sup>3</sup>College of Pharmacy, JSS Academy of Higher Education and Research, Mysore, India. E-mail: u28349505@tuks.co.za\_

Acne vulgaris is a chronic inflammatory skin disorder affecting the pilosebaceous unit. Moderate forms (comedonal) are characterised by non-inflammatory lesions including blackheads and whiteheads, whereas more severe forms present as inflammatory papules, pustules, nodules and cysts. The Gram-positive microorganism, *Cutibacterium acnes*, is the major causative pathogen in acne progression characterised by a cascade of events initiated by abnormal skin keratinisation, increased sebum production, proliferation of *C. acnes* and the release of inflammatory factors. This study aimed to investigate the potential of two southern African medicinal plants, belonging to the Asteraceae and the Rosaceae family, for the treatment of acne vulgaris. In addition, the extracts were used to synthesise gold nanoparticles to determine whether their activity could be enhanced through nanoencapsulation. The antimicrobial activity was determined using the microdilution broth assay, the synthesised nanoparticles were characterised using standard techniques (High Resolution Transmission Electron Microscopy, Fourier Transform Infrared spectroscopy, Zeta-potential and Dynamic Light Scattering), quantification of phenolic content was performed using the Folin-Cioalteau reagent and the anti-biofilm activity was performed using the crystal violet staining technique. The Asteraceae species exhibited a minimum inhibitory concentration (MIC) of 7.81 µg/mL and the Rosaceae species had an MIC of 15.625 µg/mL. The Asteraceae species nanoparticles exhibited no antimicrobial activity against *C. acnes*, showed inhibition of bacterial attachment at 20.42 µg/mL (determined using phenolic content). The Rosaceae species gold nanoparticles exhibited an MIC of 10.78 µg/mL (determined using phenolic content) which was below the extract MIC. The potent antimicrobial activity of these two southern African plant extracts against the biofilm forming *C. acnes* strain (ATCC 6919) warrants further development of these extracts as commercial active ingredients or formulations to combat acne.



## **SESSION IV: ETHNOBOTANY, BIOPROSPECTING AND PRODUCTION**

### **O17. Bioexploration of indigenous plants in sub-Sahara Africa**

James Simon. Rutgers University. E-mail: [jimsimon@sebs.rutgers.edu](mailto:jimsimon@sebs.rutgers.edu)

Sub-Sahara Africa is rich in biodiversity and a paradise to explore ethnobotanically. This presentation will present three case studies on bioexploration of selected indigenous plants. From West Africa we will focus on Griffonia, Voacanga, Shea Butter and Kinkeliba using a more traditional ethnobotanical approach. In East Africa, we employed a different model and focused on those indigenous or naturalized plants that are nutrient dense. Here, our research evaluated African Indigenous Vegetables (AIVs) and identified those that are rich sources of pro-vitamin A, iron, zinc, and other health-promoting phytochemicals. We hypothesized that by improving access, availability, affordability, the consumption of AIVs would increase as would dietary diversity, food security and health and nutrition. In addition, we quantified the nutritional and anti-nutritional aspects of major AIVs. The third case focused on identifying the useful plants of Namibia, plant collection, screening using chemical and biological assays and then chemical profiling of promising species. All work was done in concert with local partners and researchers.

## SESSION IV: ETHNOBOTANY, BIOPROSPECTING AND PRODUCTION

### INVITED TALK

#### **O18. Ethnobotany and Production of Three High-value Medicinally Active Herbs, Basil, Turmeric, and Mountain mint in North Alabama**

Srinivasa Rao Mentreddy<sup>1</sup>, Lam Duong<sup>1</sup>, Trang Pham<sup>1</sup>, Suresh Kumar<sup>1</sup>, William Setzer<sup>2</sup>, Mei Wang<sup>3</sup>, Charles Cantrell<sup>3</sup>, Suresh Mathews<sup>4</sup>, and Dattatreya Gajula<sup>5</sup>. <sup>1</sup>Biological and Environmental Sciences, Alabama A&M University, Huntsville, AL 35811; <sup>2</sup>University of Alabama in Huntsville, Huntsville, AL 35802; and <sup>3</sup>USDA-ARS-NPURL, University, MS; <sup>4</sup>Samford University, Birmingham, AL; and <sup>5</sup>Coca-Cola Company, Atlanta, GA.  
E-mail: rmentreddy@gmail.com

Herbal products are gaining popularity as health supplements in the USA. About 80% of the world's population relies on herbal remedies in one form or the other, while 25% of the pharmaceutical drugs are plant-based. Among a wide range of medicinally active plants, basil (*Ocimum* spp), turmeric (*Curcuma* spp.), and mountain mint (*Pycnanthemum virginianum*) have wide applications ranging from culinary and aromatic to medicinal and confectionery purposes. Basil, particularly holy basil (*O. tenuiflorum*), has been shown to have a myriad of medicinal properties and is known as a panacea for many ailments ranging from common colds to chronic diseases such as cancers and diabetes. Turmeric, used in Indian medicine for over 4000 years, is now gaining popularity in the US as a miracle health supplement to combat inflammatory diseases and memory loss, immune deficiency, and common colds. Mountain mint is used in the medicinal and confectionery industries. Our research using basil accessions belonging to seven species has shown their anticancer and antihyperglycaemic activity. The major components in the essential oils of five selected turmeric varieties were  $\alpha$ -phellandrene (3.7–11.8%), 1,8-cineole (2.6–11.7%),  $\alpha$ -zingiberene (0.8–12.5%),  $\beta$ -sesquiphellandrene (0.7–8.0%), ar-turmerone (6.8–32.5%),  $\alpha$ -turmerone (13.6–31.5%), and  $\beta$ -turmerone (4.8–18.4%). Several of the varieties' essential oil yields and chemical profiles are comparable with those of tropical regions. Two of the four mountain mint varieties grown in North Alabama showed consistently high yield potential. They were rich in isomenthone concentrations, which increased dramatically from 20% to 69% as the season advanced, whereas pulegone and thymol tended to decrease. The essential oil components varied with varieties and with time. One of the basil accessions studied showed antidiabetic activity comparable to commonly used medicine Acarbose. The three species with desirable essential oil composition offer potential for production as high-value medicinal crops in North Alabama.



## SESSION IV: ETHNOBOTANY, BIOPROSPECTING AND PRODUCTION

### INVITED TALK

#### **O19. Instrumental Challenges with Essential Oil Adulteration**

Prabodh Satyal. Aromatic Plant Research Center, 230N 1200E, Suite 100, Lehi, UT 84043.

About 80% of commercially available, so-called “pure and natural” essential oils are adulterated for economic profit. Not only does this create concerns for the end user, this also negatively impacts hardworking farmers and their ability to make a reliable income. Typically, adulteration of essential oils occurs through the addition of synthetic and natural compounds, those related and unrelated to the oil’s composition, in order to increase profits or meet some established requirements such as ISO. There are commonly used analytical instruments like Gas Chromatography Isotope-Ratio Mass Spectrometry (GC-IRMS), Site-Specific Natural Isotope Fractionation NMR (SNIF-NMR), EntGC-MS, and components ratio quantitation to analyze essential oil samples. GC-IR-MS is often used to detect the authenticity of the EO’s origin via the isotopic ratio measurement. However, this technique has several limitations. EntGC-MS also has specific limitations, as it is only applicable to chiral molecules and the enantiomeric ratio of chiral compounds varies from origin to origin. SNIF-NMR is only useful for small molecules (such as monoterpenes) to authenticate their origin via the Deuterium ratio, but essential oils are also composed of sesquiterpenes and diterpenes. Likewise, this method requires pure isolated compounds. Every adulterant has some kind of marker or impurity. Identifying this marker is a more effective way of detecting EO adulteration. The identification of those markers in the essential oil can provide the best solution for detecting adulteration through ordinary GC-MS. Therefore, my presentation will focus on sophisticated adulteration (undetected by ordinary lab and chemists).



## **SESSION V: IN VITRO PROPAGATION AND CONSERVATION OF VALUE-ADDED PLANTS**

### **INVITED TALK**

#### **O20. Ethnobotanical knowledge paving way for medicinal compounds and value-added products in Paulownia tree**

Nirmal Joshee. Agricultural Research Station, Fort Valley State University, Fort Valley, GA 31030. E-mail: [josheen@fvsu.edu](mailto:josheen@fvsu.edu)

Bioprospecting is a systematic and organized search for useful products derived from bioresources that can be developed for overall benefits of the society and commercialization. Paulownia is a deciduous tree which is well known for its wood quality, while its medicinal properties are just gaining attention. The genus Paulownia, belonging to monogeneric Paulowniaceae family, comprises of nine species: *P. albiphloea*, *P. australis*, *P. catalpifolia*, *P. elongata*, *P. fargesii*, *P. fortunei*, *P. kawakamii*, *P. taiwaniana*, and *P. tomentosa*. Apart from its extensive use as a high-quality wood in the industrial and agricultural areas, Paulownia has also been used as an herbal medicine in traditional Chinese medicine (TCM). The “Compendium of Materia Medica” (a Chinese Materia Medica written by Li Shizhen in 1578) mentions Paulownia bark for the treatment of hemorrhoids and worms; and the flowers for reducing swelling as well as promoting hair growth. Medical research has identified several potential applications for the genus Paulownia as an antibacterial, anti-inflammatory, thirst-quenching, diuretic, antihypertensive, hemostatic and insecticidal agent. Insecticidal properties have been linked to chemicals synthesized in glandular trichomes predominantly dispersed on the surface of leaf, flower bud, and flowers and are rich in bioactive compounds. In the last few years, various Paulownia tissues and fruits have been analyzed for phytochemical screening and bioactivity of potential compounds clinically and valuable leads have been obtained. More scientific studies using in vivo disease models are warranted to elucidate the medicinal properties and underlying mechanisms of Paulownia.

## SESSION IV: ETHNOBOTANY, BIOPROSPECTING AND PRODUCTION

CONTRIBUTED ORAL TALK

### **O21. Stevia & Turmeric, consumption can help for anti-cancer benefits, know how to grow and consume**

B. K. Biswas<sup>1</sup>, V. Seymore<sup>1,2</sup>, J. Battle<sup>1,2</sup>, K. Biswas<sup>2</sup>, B. Patel<sup>2</sup>, A. Singh<sup>2</sup>, A. Chaney<sup>1,3</sup>, N. Mullen<sup>1</sup>, and A. Roberson<sup>1</sup>. <sup>1</sup>MS Biotechnology Program, College of Agriculture, Family Sciences, and Technology, Fort Valley State University, 1005 State University Drive, Fort Valley, GA 31030, <sup>2</sup>Summer Research Students, <sup>3</sup>UC-Davis Ph.D. program; E-mail: biswasb@fvsu.edu

Two very important plants-Stevia and Turmeric can be grown in our backyards for everyday consumption used for anti-cancer benefits. Stevia (*Stevia rebaudiana* Bertoni), is a plant native to Paraguay, and its leaves produce the best natural sugar substitutes. The steviol glycoside produced in stevia leaf is anti-cariogenic, anti-diabetic, antioxidant, hypotensive, anti-hypertensive, anti-microbial, anti-inflammatory, and has anti-tumor actions. On the other hand, Turmeric, (*Curcuma longa*), is also a perennial, herbaceous plant native to the Indian subcontinent and Southeast Asia. Turmeric rhizomes are used for consumption and the active ingredient in turmeric is curcumin. Some studies suggest that curcumin in turmeric has a variety of health benefits, including fighting cancer cells. Some lab studies have found it might work against lung, breast, prostate, and colon cancers. The global number of cancer patients is predicted to increase to 47% in the next 20 years. Therefore, it is better to know how to grow and consume Stevia and Turmeric as a means of cancer prevention and therapy within the context of a healthy diet. The USA imports all stevia and turmeric from outside and mainly from China, Japan. Products grown here in the USA could help. This research study to grow and consu



## **SESSION V: IN VITRO PROPAGATION AND CONSERVATION OF VALUE-ADDED PLANTS**

### **O22. Development of efficient plant regeneration systems for grapevine precision breeding and genome editing**

S.A. Dhekney, C. Jackson, A. Junior, P. Sardaru and B. Khatabi. Department of Agriculture, Food and Resource Sciences, University of Maryland, Eastern Shore, Princess Anne, Maryland, USA. E-mail: [sdhekney@umes.edu](mailto:sdhekney@umes.edu)

Grape is among the most ancient crops cultivated worldwide and is highly valued for its ability to be converted into value-added products such as wine, juice, jelly, and jam. Several pathogens affecting grapevine can result in a significant decrease in productivity of vineyards worldwide. Genetic improvement of elite table and wine grape cultivars for disease resistance and quality traits is difficult using conventional breeding techniques. Precision breeding is defined as an approach to plant genetic improvement that transfers only specific traits among sexually compatible relatives via the relatively stable mitotic cell division pathway in order to avoid the significant genetic disruption imposed upon conventional breeding by meiosis. Recent advances in *Vitis* genome sequencing combined with optimization of grapevine regeneration protocols and gene insertion techniques have enabled the successful implementation of precision breeding technology. Grapevine somatic embryos are ideal target tissues for inserting and/or editing desired traits of interest and recovery of modified grapevines. Grape species and cultivars widely vary in their embryogenic response, which necessitates the optimization of protocols for individual cultivars. Factors influencing the production of embryogenic cultures including explant type, development stage, growth media, and culture conditions have been optimized for a large number of grape species and cultivars. The development of efficient grapevine embryogenic culture systems can enable their use in precision breeding and genome editing for rapid trait improvement.


# SESSION V: IN VITRO PROPAGATION AND CONSERVATION OF VALUE-ADDED PLANTS

## INVITED TALK

### O23. Nutrients and lighting factors in controlling medicinal plant culture

Jeffrey Adelberg. Department of Plant and Environmental Sciences, Clemson University, Clemson SC 29634. E-mail: [jadlbrg@clemson.edu](mailto:jadlbrg@clemson.edu)

Plant nutrients and light spectra are both factors that profoundly influence plant growth and development. The in vitro environment offers a highly controlled platform to explore the effects of environmental on developmental growth and phytochemistry in clonal germplasm free from confounding micro-organisms, weather, and soil conditions. In clones of turmeric, *Curcuma longa* L., high nitrate, low ammonium, medium rich in phosphate was most useful to grow rhizomes and more water was needed allow maximal use of nitrate, and more sucrose was necessary for maximal use of phosphate. The elevated phosphate allowed rhizome dry mass and vegetative propagules to develop but interfered with the formation of curcuminoids and terpenes. Bioreactors and advanced DOE experimental models allowed a small experiment to reduce many of these terms to practice. *Cannabis sativa* L., is sensitive to plant growth regulators in vitro. Propagation without PGR's is approached with monochromatic LED's providing both energy and regulatory signals to the plant. Light effects are based both on intensity and the ratios of the varied monochromatic sources. Both nutrient and light experiments are inherently multifactor, since ratios of factors are at least as important as absolute amounts. The experimental designs, in vitro methods, and special facilities needed to conduct this type of work will be shown.



## **SESSION V: IN VITRO PROPAGATION AND CONSERVATION OF VALUE-ADDED PLANTS**

### **INVITED TALK**

#### **O24. In vitro regeneration of medicinal plants and analysis of structural plasticity of cells and tissues towards changed environmental conditions: A way forward to develop climate-resilient plants**

Manokari M<sup>1,2</sup> and Mahipal S Shekhawat<sup>1</sup>. <sup>1</sup>Plant Biotechnology Unit. K. M. Government Institute for Postgraduate Studies and Research, Puducherry – 605 008, India and <sup>2</sup>Department of Botany, Siddha Clinical Research Unit, Central Council for Research in Siddha, Palayamkottai, Tamil Nadu – 627 002, India. E-mail: shekhawatms@dhtepdy.edu.in; msspd@gmail.com

The impact of climate change is evident on various ecosystems on the earth. The response of plant communities toward changed environmental conditions starts at the molecular level and is visible at cell and tissue levels (via microscopy). In vitro regeneration systems are highly useful in the mass production of genetically uniform plantlets. This technique provides controlled (simulated) environments to check how plants respond toward changed environments. The blending of microscopy with in vitro plant regeneration technology may shed light on the microstructural and histochemical developments taking place in plants under in vitro (fully heterotrophic), ex vitro (mixotrophic), and in vivo (fully autotrophic) environments. Plants that are growing under fully controlled conditions display underdeveloped foliar anatomical traits such as thin cuticle, unorganized epidermal cells, and fewer mesophyll tissue density and vascular elements with reduced deposition of lignins, cutin, pectin, tannins, and polyphenolic compounds. The mixotrophic environment impacts the improvements in the cuticle, development of uniform epidermal layer, increased mesophyll and vascular elements, and histochemical traits. The field-grown plants show well-developed dermal, ground, and vascular tissue systems along with increased deposition of primary and secondary metabolites. These gradual changes positively affect the microstructural and physiological status of plantlets to withstand abiotic stresses in the field conditions which could assist the plantlets in the tolerance to elevated temperature and light. Thus, the microscopic evaluation of plantlets could help to better understand the anatomical and histochemical process of adaptation of plants and enhanced survival of plantlets under changed environmental conditions.

**11<sup>th</sup> ANNUAL HYBRID CONFERENCE**

**American Council for Medicinally Active Plants (ACMAP)**

**EXPLORING THE UNEXPLORED: SCIENCE AND APPLICATIONS OF MEDICINAL PLANTS**

**JUNE 28 - JULY 2, 2022 | PUERTO RICO, USA**





## SESSION V: IN VITRO PROPAGATION AND CONSERVATION OF VALUE-ADDED PLANTS

### INVITED TALK

#### **O25. Conservation of Mexican medicinal species with sedative and anti-inflammatory properties**

Reinier Gesto-Borroto<sup>1</sup>, Alexandre Cardoso-Taketa<sup>1</sup>, Anabel Ortíz-Caltempa<sup>2</sup>, Gema Balderas<sup>2</sup>, Raúl E. Alcalá<sup>3</sup>, María Luisa Villarreal<sup>2</sup>. <sup>1</sup>Dirección de Ingeniería en Biotecnología, Universidad Politécnica del Estado de Morelos 62550, Jiutepec, Morelos, <sup>2</sup>México Centro de Investigación en Biotecnología Universidad Autónoma del Estado de Morelos 62209, Cuernavaca, Morelos México. <sup>3</sup>Centro de Investigación en Biodiversidad y Conservación, Universidad Autónoma del Estado de Morelos, 62209, Cuernavaca, Morelos, México

Galphimia species of the Malpighiaceae family, are used as a popular sedative and anti-inflammatory medications in Mexico. The active sedative compounds are nor-secofriedelane triterpenes known as galphimines. Using metabolic profiling and, molecular DNA barcodes, we previously demonstrated that nine populations botanically classified as *Galphimia glauca* belong to four different species of the genus *Galphimia*, and that only one species exhibited the sedative properties; however, all the collected species showed anti-inflammatory activity. In a different study, we tested the hypothesis that this accumulation responds to genetic control of *Galphimia* species. Individuals were collected from six previously studied populations, located in five Mexican states (Doctor Mora in Guanajuato, Jalpan in Querétaro, Cuernavaca and Tepoztlán in Morelos, Guadalajara in Jalisco, and Tuxtla Gutiérrez in Chiapas), and transplanted to grow under uniform greenhouse conditions, for a period of 11 months. A phytochemical analysis using chromatographic procedures was performed for all individuals upon arrival and after 11 months of controlled growth. Results indicated that only two populations (Doctor Mora, Guanajuato, and Jalpan, Querétaro), out of the six that were collected, produced galphimines before and after controlled growth, exhibiting the same chemotype pattern in the greenhouse, as in the wild. Evident changes in foliar morphology, probably due to phenotypic plasticity, were observed in all the acclimatized plants.



## SESSION V: IN VITRO PROPAGATION AND CONSERVATION OF VALUE-ADDED PLANTS

CONTRIBUTED ORAL TALK

### **O26. Machine learning models in plant tissue culture: a case study of Brahmi**

Ramana Gosukonda<sup>1</sup>, Nirmal Joshee<sup>2</sup>, Ajit Mahapatra<sup>2</sup>. <sup>1</sup>Department of Agricultural Sciences and <sup>2</sup>Agricultural Research Station, \*, College of Agriculture, Family Sciences and Technology, Fort Valley State University, Fort Valley, GA 31030. E-mail: gosukonr@fvsu.edu; josheen@fvsu.edu; mahapatraa@fvsu.edu

The objective this study is to describe applications of machine learning models in plant tissue culture using *Bacopa monnieri* (L.) as a case study. *Bacopa* is a medicinal Ayurvedic herb that has been used centuries for improving memory and cognitive skills, insomnia, anxiety, depression and seizures. These therapeutic benefits of *B. monnieri* caused an increased demand for its use in commercial preparations. Tissue culture propagation offers clean, identical, and compatible Brahmi biomass in a large scale for downstream processing. On the other hand, in vitro growth kinetics can be viewed as a multi-variable procedure impacted by the physical conditions (such as types, age, size, density, position, and source of the explants, light intensity, and temperature), the composition of culture media (macro and micronutrients, amino acids, vitamins and other nitrogen elements, organic supplements, carbon sources, agar), multiple endogenous and exogenous phytohormones, and their interactions. The combination of these factors often causes plant in vitro processes to be non-linear and non-deterministic developmental progressions. Because machine learning models such as Artificial neural networks (ANN), Random Forests (RF), Self-organizing maps (SOM) have inherent ability to handle high biological variability and the uncertainty associated with the tissue culture methods, they have proven to be appropriate approaches for modeling the non-linear and ill-defined in vitro biological processes. Lately, the use of machine learning approaches to accurately predict in vitro growth kinetics have gained importance in validating plant tissue culture processes and helping the industry to scale up productions by fine-tuning the automation process.

## SESSION V: IN VITRO PROPAGATION AND CONSERVATION OF VALUE-ADDED PLANTS

CONTRIBUTED ORAL TALK

### O27. Histological examination of in vitro hyperhydric cultures of *Scutellaria lateriflora*

Samantha H. Sherman and Nirmal Joshee. Agricultural Research Station, Fort Valley State University, Fort Valley GA 31030.

E-mail: [Samantha.Sherman@fvsu.edu](mailto:Samantha.Sherman@fvsu.edu); [josheen@fvsu.edu](mailto:josheen@fvsu.edu)

*Scutellaria lateriflora* (American skullcap) belongs to the mint family, Lamiaceae. It is a facultative wetland plant distributed across most of North America. Traditionally, American skullcap has been used for anxiolytic properties, treatment of insomnia, and inflammation and is also available in the herbal health products market as tea, capsules, extracts, and tinctures. Our previous studies on *S. lateriflora* to assess the effects of three culture systems (semi-solid agar, liquid stationary, and liquid agitated) on biomass accumulation revealed poor biomass yield coupled with high hyperhydricity in liquid stationary culture system. Two node explants were inoculated on the basal Murashige and Skoog media supplemented with sucrose (non-reducing) or maltose (reducing) at 0 %, 3 %, 5 %, 7 %, and 10 % (w/v). With increase in the carbon source concentration, plant quality decreased producing hyperhydric shoots which exhibited dark green to purple, brittle, swollen stems, and small succulent leaves. There was a relationship observed with the increase in carbon source concentration and a decrease in the production of aerial parts. On the contrary, with the increase in sugar concentration, more number of thickened roots with purple pigmentation were produced. The liquid stationary system was selected to further study morphology and anatomy of hyperhydric plants by inoculating two node explants for six weeks on basal MS medium. Leaves, stems, and roots were collected from nine treatments of carbon source for histological examination using paraffin wax sections stained with safranin and fast green. Hyperhydric tissues differed in cuticle deposition, organization of different cell layers, and presence of large intercellular spaces in comparison to control plants. To the best of our knowledge, this is the first report on the hyperhydricity in *S. lateriflora* in relation to type of culture, carbon source and concentration in an in vitro system.



## **SESSION VI:**

### **BIOACTIVE COMPOUNDS OF PLANTS AND THEIR ROLES IN HUMAN HEALTH II (INFLAMMATORY DISEASES)**

#### **O28. Flavonoids regulate macrophage immune function controlling cancer and halting obesity**

Andrea I. Doseff. Department of Physiology, Department of Pharmacology and Toxicology, Michigan State University, East Lansing, MI, 48824, USA.  
E-mail:doseffan@msu.edu

Chronic inflammatory disease, including cancer and obesity, are reaching epidemic levels worldwide, inflicting life-threatening conditions and great socioeconomic burden. Thus, there is a great need to identify new therapeutic approaches to control inflammation. Flavonoids, including the flavones, are the largest class of bioactive dietary phytochemicals attracting great interest for their beneficial effects in the treatment and prevention of chronic inflammatory diseases. We showed that the dietary flavone apigenin and a celery-based food rich in apigenin (CEBAR) reduces inflammation by modulating macrophage immune function. Using preclinical models of breast cancer and obesity, we found that apigenin decreases the number of aberrant macrophages at inflammatory sites. We found that apigenin targeting NFkB, a key transcription factor in inflammation, reduces tumor growth and metastasis. Our studies using RNAseq analyses revealed the gene regulatory networks modulated by apigenin. Extending our studies to models of obesity-induced inflammation, we showed that apigenin reducing macrophage infiltration rewires adipocytes and the microbiome resulting in weight reduction. I will discuss our results and provide a new framework to understand how medicinally active phytochemicals result in their recognized health benefits in chronic inflammatory diseases. These findings may impact the clinical use of flavones in the prevention and treatment of chronic inflammatory diseases.

## **SESSION VI:**

# **BIOACTIVE COMPOUNDS OF PLANTS AND THEIR ROLES IN HUMAN HEALTH II (INFLAMMATORY DISEASES)**

### **INVITED TALK**

#### **O29. Anti-inflammatory Interactions of Tannins with the Intestinal Microbiome**

Susanne U. Mertens-Talcott <sup>‡</sup> Hyemee Kim <sup>‡</sup> Maria Joselyn Castellon Chicas <sup>‡</sup>  
Stephen T. Talcott <sup>2</sup>. <sup>1</sup>Department of Food Science and Nutrition, Pusan  
National University, Busan, 46241, South Korea. <sup>2</sup>Department of Food Science  
and Technology, Texas A&M University, College Station, TX, 77843, USA  
Email: SMTalcott@tamu.edu

The global botanical market is projected to reach USD 49.36 billion by 2027, many based on a polyphenolic bioactive in the prevention of a wide array of ailments, including intestinal inflammation. Emerging evidence suggests that inter-individual variability of polyphenol metabolism has contributed significantly to inconclusive clinical trials with polyphenolics, and this is likely to be significantly mediated by differences in the intestinal microbiota. Gallotannins (GT) are gallic acid polymers that represent a significant portion of polyphenol intake from many fruits, green tea, spices, nuts, and botanical supplements and possess anti-inflammatory activities via IGF1R/AMPK signaling. Metabolism of GT generates the bioavailable anti-inflammatory metabolites gallic acid (GA) and pyrogallol (PG) but pharmacokinetics of these metabolites show large variation between individuals. Prevalent dysbiosis with chronic intestinal inflammatory conditions is expected to limit polyphenol metabolism, and might be ameliorated via supplementation with targeted probiotics. Two well-defined intestinal microbial enzymatic steps yield the major metabolites of GTs: a) tannase (tannin acyl hydrolase, EC 3.1.1.20) hydrolyzes galloyl-ester bonds of GT to produce GA, and b) GA-decarboxylase (EC 4.1.1.59) produces PG and both enzymes are active in probiotic strains. This presentation will present data on the anti-inflammatory interactions of probiotics and large molecular tannins relevant to intestinal inflammation.



## SESSION VI: BIOACTIVE COMPOUNDS OF PLANTS AND THEIR ROLES IN HUMAN HEALTH II (INFLAMMATORY DISEASES)

### INVITED TALK

#### **O30. Multi-omics approach reveals biomarkers of metabolic resilience in healthy participants supplemented with grape polyphenols.**

Diana E. Roopchand. Department of Food Science, NJ Institute for Food Nutrition and Health (Rutgers Center for Lipid Research and Center for Nutrition Microbiome and Health), Rutgers, The State University of New Jersey, New Brunswick, NJ, USA Email: roopchand@sebs.rutgers.edu

Grape polyphenols extracted from grape pomace are rich in proanthocyanidins (PACs) compounds which are not bioavailable; however, evidence suggests their health benefits are likely mediated via the gut microbiota and/or bacterial metabolites. Using mice fed high-fat diet (HFD) or having leptin receptor mutation (db/db), we previously showed that GP supplementation can improve symptoms of metabolic syndrome and type-2 diabetes. In particular, GP supplemented mice showed improved oral glucose tolerance in association with marked changes in gut microbial community structure, alterations in bile acid profiles, decreased intestinal FXR signaling, and reduced intestinal and hepatic markers of inflammation. To investigate the effects of GPs on the human gut microbiome and metabolome, twenty-nine healthy participants completed a single-arm longitudinal study. Serum, fecal, and/or urine samples were collected before and after 2, 4, and 10 days of GP supplementation followed by shotgun metagenomic sequencing, mass spectrometry-based metaproteomics, and/or targeted metabolomics (i.e., bile acids and polyphenols metabolites). Notable GP-induced bile acid changes appeared to be gender-specific and correlational analyses uncovered novel associations between gut microbes and metabolites associated with metabolic resilience.

## SESSION VI: BIOACTIVE COMPOUNDS OF PLANTS AND THEIR ROLES IN HUMAN HEALTH II (INFLAMMATORY DISEASES)

### INVITED TALK

#### **O31. Metabolic benefits of grape polyphenol consumption on bile acid signaling in mice with intestine- and liver-specific deletion of Fxr**

Kevin M. Tveter<sup>1</sup>, Sriya Sadangi<sup>3</sup>, Karen Bacalia<sup>2</sup>, Yue Wu<sup>1</sup>, Hung Hoang<sup>3</sup>, Esther Mezhibovsky<sup>2</sup>, Ke Sui<sup>1</sup>, and Diana E. Roopchand<sup>1</sup>. <sup>1</sup>Rutgers, The State University of New Jersey, Department of Food Science, Institute for Food Nutrition and Health, 61 Dudley Road, New Brunswick, NJ 08901, USA. <sup>2</sup>Rutgers, The State University of New Jersey, Department of Nutritional Sciences, Institute for Food Nutrition and Health, 61 Dudley Road, New Brunswick, NJ 08901, USA. <sup>3</sup> Rutgers, The State University of New Jersey, Department of Cell Biology and Neuroscience, New Brunswick, NJ 08901, USA. E-mail: roopchand@sebs.rutgers.edu, Kevin.tveter@rutgers.edu

Bile acids (BA) are a unique class of lipids pivotal for digestion, metabolism, and absorption of other lipid species. Primary BAs (PBA) are produced using cholesterol in the liver and released into the gastrointestinal tract where they are modified by gut microbes into secondary BAs (SBA). Evidence suggests the gut microbiome may regulate energy metabolism by altering BA which, in-turn, may modulate energy and lipid metabolism via their differential affinity for BA receptors like nuclear transcription factor, Farnesoid X receptor (Fxr). Proanthocyanidins (PACs) are a class of polyphenols found in fruits that epidemiological and clinical evidence suggests may be beneficial, when consumed, for the prevention and management of cardiometabolic diseases. We have previously seen that consumption of a PAC-rich, grape polyphenols (GP) extract dramatically remodeled the gut microbial communities and BA profile of diabetic (db/db) mice to inhibit intestinal Fxr signaling and improve glucose metabolism. To investigate the tissue-specific requirement of Fxr in mediating the glycemic benefits of GP, we bred mice using Cre-Lox recombinase technology to breed mice with intestine and liver specific deletion of Fxr. Mice with liver- (AKO) and intestine (VKO)-specific deletion of Fxr and wild-type (WT) floxed littermates were fed high-fat diet (HFD) or HFD supplemented with 0.5% GP (HFD-GP) for 12 weeks. Metabolic phenotyping, BA profiles and gene expression of metabolic tissues, and cecal gut microbial composition were analyzed. Consumption of GPs did not have a impact on the metabolic phenotype of VKO mice as these mice are resilient to HFD-induced obesity. Interestingly, WT and AKO mice fed HFD-GP had decreased body weight and improved glucose metabolism compared to HFD-fed counterpart; however, BA profiles of liver and serum were markedly different. In all three strains, GPs significantly decreased expression of inflammatory markers (Tnfa, IFN $\gamma$ , Nos2) in one or more metabolic tissues

### **11<sup>th</sup> ANNUAL HYBRID CONFERENCE**

**American Council for Medicinally Active Plants (ACMAP)**

**EXPLORING THE UNEXPLORED: SCIENCE AND APPLICATIONS OF MEDICINAL PLANTS**

**JUNE 28 - JULY 2, 2022 | PUERTO RICO, USA**



## SESSION VI: BIOACTIVE COMPOUNDS OF PLANTS AND THEIR ROLES IN HUMAN HEALTH II (INFLAMMATORY DISEASES)

### INVITED TALK

#### **O32. Flavone apigenin regulates innate immune response through RNA binding protein hnRNPA2**

Meenakshi Sudhakaran <sup>1,2</sup>, Daniel Arango<sup>4</sup>, Andrea I. Doseff <sup>2,3</sup>. <sup>1</sup>Molecular, Cellular and Integrative Physiology Program, <sup>2</sup>Department of Physiology, <sup>3</sup>Department of Pharmacology and Toxicology, Michigan State University, MI. <sup>4</sup>Department of Pharmacology, Northwestern University, IL.  
E-mail: sudhaka7@msu.edu; doseffan@msu.edu

Inflammatory diseases such as cancer are characterized by an abnormal expansion of specific myeloid cells, including monocyte and macrophages that fuel tumor growth and dysregulated immune response. Thus, there is a need to identify new approaches to restore immune homeostasis. Apigenin, a naturally abundant flavone, is emerging as a potential immuno-regulatory compound. Apigenin inhibited pro-inflammatory cytokine production in LPS-stimulated monocytes and in vivo transgenic mice. To investigate the mechanisms responsible for its function, we identified direct targets of apigenin using a phage-display library coupled with next-generation sequencing (PD-Seq). Among them, the RNA binding protein hnRNPA2 was found as a high affinity target. While the role of hnRNPA2 in cancer growth is well recognized, its function in inflammation has not been investigated. In this study we reveal that apigenin through hnRNPA2 modulates inflammation. We found that apigenin diminishes LPS- and TNF $\alpha$ -stimulated inflammation through hnRNPA2. Lack of hnRNPA2 inhibits NF- $\kappa$ B, a transcription factor crucial in inflammation, as shown by reporter assays in stimulated macrophages. Deletion of hnRNPA2 using siRNA, similar to apigenin, inhibits the phosphorylation of NF- $\kappa$ B subunit p65 required for its transcriptional activity. Inhibition of NF- $\kappa$ B by apigenin or lack of hnRNPA2 resulted in suppression of pro-inflammatory cytokine expression in stimulated macrophages. Immunoprecipitation assays showed that hnRNPA2 associates with NF- $\kappa$ B. Genome-wide expression of stimulated macrophages suggests an hnRNPA2-dependent regulatory network contributing to NF- $\kappa$ B regulation. Together, our study identified a new role of how apigenin through binding with hnRNPA2 modulates NF- $\kappa$ B-dependent inflammation, thereby highlighting the relevance of apigenin as a potential immune-modulatory agent to prevent and treat inflammatory diseases.

## SESSION VII: HEMP AND MEDICAL CANNABIS

### O33. Genetic complexity in Industrial hemp germplasm

Aruna D Weerasooriya. College of Agriculture & Human Sciences, Cooperative Agricultural Research Center, Prairie View A&M University, Prairie View, Texas 77446  
E-mail: [adweerasooriya@pvamu.edu](mailto:adweerasooriya@pvamu.edu)

Industrial hemp (*Cannabis sativa* L.) is a multifaceted yet controversial crop that is long associated with humans. It contains genotypes of both non-narcotic hemp that has numerous industrial uses and intoxicating marijuana where their identities are mainly based on a threshold of THC (delta-9-tetrahydrocannabinol) concentration. Industrial hemp exhibits a worldwide distribution and is considered a lucrative industrial crop. The recent legalization of hemp provided a major economic impact on US agriculture. Therefore, elite hemp strains for different agroecological conditions and end-usages (e.g., fiber, seed grain, phytochemical, and other value-added products) should be identified and introduced to hemp breeding programs. We have acquired hemp genetic resources through exchange and exploration programs and cultivated them in controlled environmental conditions to obtain the best quality research samples to identify molecular signatures of industrial hemp using phenomic, metabolomic, and genomic approaches and test their agronomic performance in field cultivations. A collection of well-curated herbarium specimens of all cultivated strains was established and a digital repository was created. The superior strains resulting from this intensive filtering criteria will be identified for inclusion in the USDA Hemp Germplasm Repository at Cornell AgriTech in Geneva, New York, and also to use in field cultivations, breeding, and bioengineering experiments.



## SESSION VII: HEMP AND MEDICAL CANNABIS

### INVITED TALK

#### 034. Endocannabinoid receptor pharmacology of rare cannabinoids

Gia-Nam Ngyuen <sup>1</sup>, Erin Jordan <sup>1</sup>, Oliver Kayser <sup>1</sup>. <sup>1</sup>TU Dortmund University, Technical Biochemistry, Emil-Figge-Strasse 66, 44227 Dortmund, Germany.  
E-mail: [oliver.kayser@tu-dortmund.de](mailto:oliver.kayser@tu-dortmund.de)

Cannabinoids are important terpenophenolics in *Cannabis sativa* L. and are found at high concentrations in trichomes. Today, most cannabinoids tested have been isolated from the cannabis plant for their activity on endocannabinoid receptors CB1 and CB2. Unfortunately, most of the rare cannabinoids are not easily accessible why a limited number has been tested so far. In our continued work we have synthesized a series of cannabinoids with varying aliphatic chain lengths (C5, C3, C1, C0). Efficient syntheses of eight key cannabinoids were established and optimized. Predominant cannabinoids like cannabigerol (CBG-C) and cannabidiol (CBD<sub>5</sub>-C) were prepared from olivetol via regioselective condensation. Further treatments of CBD led to  $\Delta^9$ -tetrahydrocannabinol (THC-C),  $\Delta^8$ -iso-tetrahydrocannabinol ( $\Delta^8$ -o-THC-C) and cannabinol (CBN<sub>5</sub>-C). Alternatively, a [3+3] annulation between olivetol and citral yielded the minor cannabinoid cannabichromene (CBC-C), which was converted into two very rare polycycles cannabicyclol (CBL-C) and cannabicitran (CBT-C) in an one-pot reaction (Fig.1). Finally, all eight syntheses were extended by utilizing resorcinol and two phenolic analogs, achieving a cannabinoid group with more than 30 compounds through a facile synthesis strategy. All cannabinoids were tested for recombinant CB1, CB2, FAAS and MAGL activity or inhibition. Based on the different analogs of tested cannabinoids surprising direct and indirect effects were observed for cannabinoids with varying chain lengths. In this talk, we provide a structure-activity-relationship (SAR) and explain the potential interactions of cannabinoids with protein domains of endocannabinoid receptors.

## SESSION VII: HEMP AND MEDICAL CANNABIS

### INVITED TALK

#### **O35. Optimizing propagation techniques and management practices for Cannabis sativa L. production in the U.S. Mid-Atlantic region**

S.A. Dhekney, E. Lindsay, A. Junior, C. Jackson, P. Sardaru, M.H. Foland and B. Khatabi. Department of Agriculture, Food and Resource Sciences, University of Maryland Eastern Shore, Princess Anne, Maryland 21853, USA.

E-mail: [sdhekney@umes.edu](mailto:sdhekney@umes.edu)

The legalization of industrial hemp under the 2014 and 2018 US farm bills has led to a rapid increase in hemp production in the United States. Legally, hemp is classified as any cultivar of Cannabis sativa that has a delta 9 THC (tetrahydrocannabinol) level of 0.3% and such cultivars have been federally mandated for seed (oil, food and feed), fiber and cannabinoid production. Hemp has diverse applications including its use in food, fiber, and medicinal products. Hemp propagation for cannabinoid production is achieved using feminized seed or softwood cuttings. Feminized seed production involves treating female hemp plants with plant growth regulators to induce sex reversal, leading to the development of male flowers and self-fertilization. Seed and seedlings produced from treated plants are predominantly female thereby eliminating the need for clonal propagation. The quality of hemp planting material obtained through asexual propagation and feminized seed is often compromised due to the transmission of pathogens. Large-scale propagation of uniform, disease-free female plants can be achieved through in vitro culture techniques such as micropropagation. Hemp growth and cannabinoid levels is significantly influenced by genotype, soil and environmental conditions, which necessitates identification of suitable cultivars for specific macroclimates. We have optimized protocols for feminized seed production and micropropagation, for large-scale propagation of elite hemp clonal material. We are also evaluating several hemp cultivars for their suitability in cannabinoid production under indoor and field conditions. The optimization of propagation techniques and cultivar screening trials will enable growers select the appropriate legal cultivars for maximum biomass yield and cannabinoid levels and boost expansion of the hemp industry in the United States.

### **11<sup>th</sup> ANNUAL HYBRID CONFERENCE**

**American Council for Medicinally Active Plants (ACMAP)**

**EXPLORING THE UNEXPLORED: SCIENCE AND APPLICATIONS OF MEDICINAL PLANTS**

**JUNE 28 - JULY 2, 2022 | PUERTO RICO, USA**



## SESSION VII: HEMP AND MEDICAL CANNABIS

### INVITED TALK

#### O36. Chasing the entourage effect in Cannabis

Wesley M. Raup-Konsavage<sup>1</sup>, Nurgul Carkaci-Salli<sup>1</sup>, Diana E. Sepulveda<sup>1,2</sup>, Vengadeshprabhu Karuppa Gounder<sup>3</sup>, Fadia Kamal<sup>3</sup>, Nicholas M. Graziane<sup>1,2</sup>, Kent E. Vrana<sup>1</sup>. <sup>1</sup>Department of Pharmacology, <sup>2</sup>Department of Anesthesiology & Perioperative Medicine, <sup>3</sup>Department of Orthopedics and Rehabilitation, Penn State University College of Medicine, Hershey PA 17033.  
E-mail [wkonsavage@pennstatehealth.psu.edu](mailto:wkonsavage@pennstatehealth.psu.edu)

Cannabis produces a large number of compounds with potential medical uses, these compounds include cannabinoids, terpenes, and flavonoids. It is speculated that synergism between these three groups of molecules, known as the “entourage” effect, substantiates the use of the whole plant for medicinal purposes. However, pure compounds are typically the standard for treatments in the medical and pharmaceutical fields. To examine the effectiveness of the “entourage effect”, we compared low-THC containing cannabis (hemp) extracts with pure cannabinoids in several disease models. We found that pure cannabidiol (CBD) has a modest but significant impact on colorectal cancer, glioblastoma, and melanoma cell viability. A comparison of three commercial CBD oils demonstrated markedly different effects on cell viability, however, no oil was superior to pure CBD, the IC<sub>50</sub> values for pure CBD were lower than those for the most potent oil. This argues against the existence of an “entourage” effect in potential cancer treatments. In a murine model of chemotherapy-induced peripheral neuropathy (CIPN), we found that cannabigerol (CBG) significantly reduced hyperalgesia, measured using the von Frey test. Additionally, we found that CBG/CBD oil (at a comparable CBG concentration) was significantly more effective at reducing hyperalgesia than pure CBG in mice expressing CIPN. Furthermore, in a murine model of osteoarthritis, we found that CBG/CBD oil and pure CBG were both similarly able to reduce pain and disease progression; however, unlike CBG/CBD oil, pure CBG did not produce anti-inflammatory activity. These data suggest that there is something in the oil that contributes to the antinociceptive properties of these cannabinoids, additional studies are currently underway. In conclusion, if an “entourage” effect exists it appears to be dependent upon the condition being treated and not a more universal property of cannabis compounds.

**11<sup>th</sup> ANNUAL HYBRID CONFERENCE**

**American Council for Medicinally Active Plants (ACMAP)**

**EXPLORING THE UNEXPLORED: SCIENCE AND APPLICATIONS OF MEDICINAL PLANTS**

**JUNE 28 - JULY 2, 2022 | PUERTO RICO, USA**



## **SESSION VII: HEMP AND MEDICAL CANNABIS**

### **CONTRIBUTED ORAL TALK**

#### **O37. Cannabis testing labs; Quality control and standardization**

Niranjan Aryal. Lab Director at VRX Labs, Long Beach California.  
E-mail: naryal101@gmail.com

Cannabis testing laboratories play a pivotal role in the multidimensional cannabis industry. Production companies to the consumers, whole cannabis community rely on the analytical results produced by the laboratories. Given the immense importance, it is the responsibility of testing laboratories to produce precise, accurate, and consistent results for every sample they analyze. Several uncertainty factors including the Quality control (QC) checks, that a laboratory runs in every batch, are determining factors for the certainty of the results. Quality control checks are not only to satisfy the regulatory boards but to have confidence in the results produced. From calibration to the sample runs, every batch should consist of quality control checks. It should be ensured that QC checks such as method blank, technical blank, laboratory control sample, and continuing calibration verifications are passed before analyzing the data for the rest of the samples. Blind testing such as timely proficiency testing helps in maintaining the accuracy of the instruments and the analysis. When the whole cannabis industry is proceeding towards the standardization, testing laboratories as well are working on it. As of now, a clearer picture for the ultimate standardization of Cannabis testing laboratories is not visible, however, the debate has been started on how the standardization should look like, a plateau, or a peak? When several countries are preparing to legalize cannabis and be part of the international market, a worldwide accepted standardization on testing is much awaited. I will be presenting on how the quality control in testing laboratories impacts the whole cannabis industry and how a testing laboratory can maintain a good quality control system. I will also be discussing how the future looks like for testing laboratories.

## SESSION VII: HEMP AND MEDICAL CANNABIS

### CONTRIBUTED ORAL TALK

#### **O38. Cannabidiol improves inflammation, glucose metabolism and bone loss in a murine model of female post-menopause in association with a bloom of *Lactobacillus* sp.**

Ke Sui<sup>1</sup>, Kevin Tveter<sup>1</sup>, Fiona Gaile Bawagan<sup>1</sup>, Patricia Buckendahl<sup>2</sup>, Savannah Martinez<sup>3</sup>, ZehraJaffri<sup>3</sup>, Avery Macdonell<sup>3</sup>, Rocio Duran<sup>1</sup>, Sue Shapsess<sup>4</sup>, and Diana E. Roopchand<sup>1\*</sup>. <sup>1</sup>Department of Food Science, NJ Institute for Food Nutrition and Health (NJIFNH, Rutgers Center for Lipid Research and Center for Nutrition Microbiome and Health), <sup>2</sup>Molecular Imaging Center, <sup>3</sup>Department of Biological Sciences, <sup>4</sup>Department of Nutritional Science, NJIFNH (Center for Human Health and Performance), Rutgers, The State University of New Jersey, New Brunswick, NJ, USA.

E-mail: ke.sui@rutgers.edu; roopchand@sebs.rutgers.edu

The predominant form of estrogen, 17 $\beta$ -estradiol, declines during menopause resulting in increased risk of gastrointestinal disorders, cardiometabolic disease, and osteoporosis. Hormone replacement therapy can alleviate menopausal symptoms but also increases the risk of cancer and heart disease. Cannabidiol (CBD) has been reported to have anti-inflammatory effects therefore we investigated whether it could improve the consequences of estrogen deficiency in ovariectomized (OVX) C57BL6/J female mice. At 12 weeks of age mice underwent ovariectomy (OVX, n= 24) or sham surgery (SS, n= 24) followed by peroral administration of CBD (25 mg/kg) or vehicle (VEH, sesame oil) for 18 weeks. CBD treatment increased oxygen consumption and energy expenditure in OVX mice. In OVX mice, CBD treatment improved oral glucose tolerance compared to VEH-treated mice. CBD-treated OVX group had lower colonic gene expression of IL-6 and TNF $\alpha$  and increased gene expression level of tight junction protein ZO-1 compared to VEH-treated mice. Dual-energy X-ray absorptiometry showed that compared to VEH-treated OVX mice, CBD-treated OVX mice had increased whole body and femur bone mineral density. High resolution micro-computed tomography showed that CBD treatment improved trabecular thickness and bone volume/trabecular volume. Compared to the VEH-treated OVX group, changes of femoral mRNA expression of IL-6, RANKL and CB2 of CBD-treated OVX group indicated reduced inflammation and bone loss. 16S rRNA amplicon sequencing analysis revealed the fecal microbiota of CBD-treated OVX mice had increased relative abundance of *Lactobacillus*, which is associated with reduced bone loss and inflammation. In CBD-treated OVX mice, bile acid membrane receptor (Tgr5) in femur was activated and increased serum level of tauroursodeoxycholic acid (TUDCA) was observed. Overall, these data suggest that CBD may be useful in post-menopause for alleviating intestinal inflammation and bone loss as well as improving glucose metabolism via alteration of gut microbiome like increased relative abundance of *Lactobacillus* species.

### **11<sup>th</sup> ANNUAL HYBRID CONFERENCE**

#### **American Council for Medicinally Active Plants (ACMAP)**

EXPLORING THE UNEXPLORED: SCIENCE AND APPLICATIONS OF MEDICINAL PLANTS

JUNE 28 - JULY 2, 2022 | PUERTO RICO, USA

## SESSION VIII: PLANT AND ALGAE DERIVED COSMECEUTICALS AND NUTRACEUTICALS

### 039. Reinforce urban skin by connecting with the origins

Namrita Lall<sup>1,2,3</sup>. <sup>1</sup>Professor and a Research Chair; Medicinal Plant Science, University of Pretoria, South Africa, <sup>2</sup>School of Natural Resources, University of Missouri, Columbia, USA, <sup>3</sup>College of Pharmacy, JSS Academy of Higher Education and Research, Mysore, India. E-mail: namrita.lall@up.ac.za

Cosmeceuticals are hybrids between drugs and cosmetic products and are able to enhance both health and beauty by external application which is very important for women specially. With about 25 000 known species, South Africa, is third only to Brazil and Indonesia as far as biodiversity is concerned. This constitutes about one tenth of all plant species in the world. Whether searching for ways to treat serious diseases like cancer and tuberculosis (TB), or formulating new acne creams and toothpastes that fight gum disease, one can be convinced that solutions can be found in South Africa's indigenous plants. The vast traditional knowledge about SA's plants is still untapped, and there is a huge amount of work needed to verify if and how traditional remedies and local plants actually work.

The synergy of usage of plants traditionally coupled with all the rare plants found in this country prompted us to evaluate the potential of unexplored plants and develop products which can be applied topically. A number of plants have been proven to have effect for skin- problems for example melasma, spots, pigmentation, acne etc. and effective cosmeceutical-prototypes products have been prepared from those. Not all work; what are bring to the lab, on average only 2% reach a stage which are subjected to clinical studies. Due to the already existing products in the cosmetic market, one needs to check carefully how well the efficacy of newly invented SA indigenous plants; compares with the available ones.

The purpose of this talk is to provide a holistic overview on the application of medicinal plants for the disorders of the skin including pigmentary disorders, wrinkle problems and melanoma. Once the medicinal products are developed, the dynamics of various ways for benefitting the people who have been using plants for centuries, will be discussed in the talk.



## **SESSION VIII:**

# **PLANT AND ALGAE DERIVED COSMECEUTICALS AND NUTRACEUTICALS**

### **INVITED TALK**

#### **O40. Health-promoting effects of functional foods**

Sun-Ok Lee, Cindi Brownmiller, and Inah Gu. Department of Food Sciences, University of Arkansas, Fayetteville, AR 72704, USA. E-mail: [sunok@uark.edu](mailto:sunok@uark.edu)

Phytochemicals are not essential compounds that are usually present in small quantities as natural constituents in food. However, they have received scientific attention due to their protective effect against a number of chronic diseases including heart disease, diabetes, various type of cancer, stroke, and metabolic syndrome. Public interest in foods that provide health benefits beyond basic nutrition has increased dramatically due to several factors, including an increase in healthcare costs and the desire to improve personal health. Current research efforts are focused not only on the development of health-promoting foods for health maintenance throughout the life cycle, but also on researching phytochemicals to evaluate the necessary dosage for optimum health. The ability of a compound to exert its biological effects depends on its bioavailability dose, rather than the administrated dose. Our recent data have shown the health benefits or bioavailability of phytochemicals in berries (blackberry, red raspberry, and blueberry), watermelon (flesh, rind, and skin), sweet potato leaves, and onion peels (byproducts of onion).

# SESSION VIII: PLANT AND ALGAE DERIVED COSMECEUTICALS AND NUTRACEUTICALS

## INVITED TALK

### **O41. Upcycling food process byproducts for cosmeceutical applications: Case of cherry waste**

Ivan Lysenko <sup>1</sup>, Meenakshi Sudhakaran <sup>2</sup>, Hannah Decot <sup>2</sup>, Andrea I. Doseff <sup>2</sup>, Bahar Aliakbarian<sup>1,3</sup>. <sup>1</sup>The Axia Institute, <sup>2</sup>Dept. of Pharmacology&Toxicology, <sup>3</sup>Dept. of Biosystems&Agricultural Eng, Michigan State University. E-mail: bahara@msu.edu

The use of plant-derived natural antioxidants in cosmetics is preferred over synthetic antioxidants. Extracts of plant-derived antioxidants generally contain a mixture of natural compounds, which could have synergetic effects; therefore, they can have better effects and less toxicity. The tart cherry process results in the generation of byproducts such as cherry pomace (15-28% of the initial fruit) and the cherry pit (7-15% of the whole fruit), depending on the process. Currently, byproducts from cherry industries are being used as animal feed, fuel, or preparation of activated carbons, despite their nutritional potential. This project aimed to define the target components and evaluate the antioxidant activity of natural extracts from the cherry pit for cosmetic applications using cellular models. Bioactive compounds from cherry pits were extracted using a high-pressure and temperature reactor and water: ethanol (50:50 v/v). Chemical characterization of the extracts demonstrated high amounts of total polyphenols (17 mgGAE/gDry Pit), flavonoids (12 mgQUE/gDry Pit), and antiradical power (7 gDPPH/Lextract). Extracts were stable for up to 40 days when stored at 4°C. This is the first time that a food byproduct (e.g., cherry pit) has been used for cosmetic applications. Cosmeceuticals with the potential to protect the skin from inflammation injury triggered by environmental stress offer an untapped market of exponentially growing interest. The dual purpose of this project makes it different from the current natural antioxidant available in the market.



## SESSION VIII: PLANT AND ALGAE DERIVED COSMECEUTICALS AND NUTRACEUTICALS

### INVITED TALK

#### **O42. Nutraceutical and cosmeceutical properties of sericea lespedeza (*Lespedeza cuneata*)**

Thomas H. Terrill<sup>1</sup>, Nirmal Joshee <sup>1</sup>, and Andres Pech-Cervantes <sup>1</sup>.

<sup>1</sup>Agricultural Research Station, Fort Valley State University, Fort Valley, GA 31030. E-mail: terrillt@fvsu.edu

*Sericea lespedeza* (SL; *Lespedeza cuneata*), a warm-season perennial legume native to Asia and Eastern Australia, has been used in the United States for over 100 years for soil conservation and as an inexpensive hay and grazing crop. Interest in the plant as a nutraceutical forage has surged over the past 20 years due to its excellent bioactivity against gastrointestinal nematodes and coccidia in livestock and its ability to reduce methane emissions from ruminant animals. While much of this bioactivity has been attributed to the high concentration and unique structure of the condensed tannins in SL, which consist of up to 98% prodelphinidin subunits, recent research has revealed the presence of a number of additional bioactive secondary compounds in the plant, including  $\beta$ -sitosterol, quercetin, kaempferol, pinitol, avicularin, juglanin, trifolin, and flavone glycosides. Due to the presence of these and other bioactive compounds, SL has been reported to possess anti-inflammatory, anti-oxidant, and anti-cancer properties, including against breast and colon cancer cells, in humans. Additional studies have demonstrated the medicinal properties of SL in treating kidney and lung disorders, as well as its anti-diabetic and hepatoprotective properties. In addition, documented anti-wrinkle, anti-melanogenic, and anti-aging properties have shown the potential value of SL for development of cosmeceutical ingredients. In addition to SL, there are several other *Lespedeza* species native to North America that may have even greater bioactivity potential than SL, but have not been adequately evaluated. Additional research is needed to unlock the nutraceutical and cosmeceutical potential of these *Lespedeza* species.

## **SESSION VIII: PLANT AND ALGAE DERIVED COSMECEUTICALS AND NUTRACEUTICALS**

### **CONTRIBUTED ORAL TALK**

#### **O43. *Euclea natalensis* A. DC., as an adjuvant, immunomodulator and hepatoprotectant**

Anna-Mari Reid<sup>1</sup>, Carel Oosthuizen<sup>2</sup> and Namrita Lall<sup>1,3,4</sup>. <sup>1</sup>Department of Plant and Soil Sciences, University of Pretoria, Pretoria, South Africa. <sup>2</sup>Drug Discovery and Development Centre (H3-D), University of Cape Town, Cape Town, South Africa. <sup>3</sup>School of Natural Resources, University of Missouri, Columbia, Missouri 65211, United States, College of Pharmacy. <sup>4</sup>JSS Academy of Higher Education and Research, Mysuru 570015, India. E-mail: u29086354@up.ac.za; namrita.lall@up.ac.za

Currently, conventional therapy is not effective against cases of infection with drug-resistant tuberculosis and severe non-compliance is hampering total eradication. Adjuvants that can be taken in conjunction with conventional treatment are of crucial importance. *Euclea natalensis* A. DC. (EN), has traditionally been used for the treatment of associated symptoms of tuberculosis, coughs and chest complaints amongst others. The root extracts of EN were shown to be very active against *Mycobacterium tuberculosis*, in-vitro, with a minimum inhibitory concentration (MIC) of 0.8 µg/ml. Due to sustainability concerns, ethanolic shoot extracts (MIC- 125 µg/ml) have been used for further development by evaluation of the immunomodulatory and hepatoprotective properties. Several further investigations into the myriad of properties of EN have been investigated such as cytotoxicity, pharmacokinetics (inhibition/stimulation of major CYP P450 enzymes), nutritional content, heavy metal analysis, microbial content, antimutagenicity and antifibrotic activity. Future projects include the PLGA nanoparticle formulation of the main active ingredient, a polyherbal formulation with several other indigenous plants and propagation trials. Development of EN has included community development, capacity building, the issuing of a bioprospecting permit and a PCT patent. Various in-vitro and preclinical in-vivo investigations have since concluded that EN is an effective adjuvant that can be used in conjunction with conventional treatment. A current licensee and proudly South African company is in the process of launching a liver protectant product with EN as the main ingredient. This product will potentially be distributed and sold at local pharmacies nationwide.

**11<sup>th</sup> ANNUAL HYBRID CONFERENCE**

**American Council for Medicinally Active Plants (ACMAP)**

**EXPLORING THE UNEXPLORED: SCIENCE AND APPLICATIONS OF MEDICINAL PLANTS**

**JUNE 28 - JULY 2, 2022 | PUERTO RICO, USA**



## **SESSION VIII: PLANT AND ALGAE DERIVED COSMECEUTICALS AND NUTRACEUTICALS**

CONTRIBUTED ORAL TALK

### **O44. Prenylated stilbenoids from peanut increase the anticancer efficacy of paclitaxel in triple-negative breast cancer cells**

Sepideh Mohammadhosseinpour<sup>1,2</sup>, Alexx Weaver<sup>1</sup>, Linh-Chi Ho<sup>1</sup>, Fabricio Medina-Bolivar<sup>1,3</sup>. <sup>1</sup>Arkansas Biosciences Institute, <sup>2</sup>Molecular Biosciences Graduate Program, <sup>3</sup>Department of Biological Sciences, Arkansas State University, Jonesboro, AR 72401. E-mails: [sepideh.mohammad@astate.edu](mailto:sepideh.mohammad@astate.edu); [fmedinabolivar@astate.edu](mailto:fmedinabolivar@astate.edu)

Triple-negative breast cancer (TNBC) is one of the deadliest forms of breast cancer. Investigating alternative therapies to increase survival rates for this disease is essential. One approach to address this issue is adjuvant therapy, where a drug is combined with a chemotherapy drug to increase its efficacy. This study aimed to examine if prenylated stilbenoids from peanut can act as adjuvants for paclitaxel, a chemotherapeutic drug with severe side effects. The prenylated stilbenoids arachidin-1 (A-1) and arachidin-3 (A-3) were produced in hairy root cultures of peanut. The cytotoxicity activity of A-1, A-3, and resveratrol (RES) alone or combined with paclitaxel was studied by checkerboard assays in TNBC cell lines MDA-MB-231 and MDA-MB-436. The apoptotic effects of this combination treatment were studied by western blotting targeting protein expression levels of PARP, caspase-8, caspase-9, and surviving and by the Apo-ONE Homogeneous Caspase-3/7 assay. To further investigate the apoptosis and cell cycle stages, cells treated with prenylated stilbenoids or RES were studied using flow cytometry. After 24 hours of treatment, A-1 exhibited higher cytotoxicity than A-3 and RES with approximately 11-fold and 6-fold lower IC<sub>50</sub>, respectively, in MDA-MB-231 cells, and 9-fold and 8-fold lower IC<sub>50</sub>, respectively, in MDA-MB-436 cells. Importantly, A-1 did not show significant cytotoxicity in the non-cancerous cell line MCF-10A. The results showed an increase in the apoptotic efficacy of paclitaxel when combined with prenylated stilbenoids. This suggests that the prenylated stilbenoid A-1 should be further explored as a potential adjuvant for chemotherapy drugs such as paclitaxel in TNBC treatment.

## **SESSION VIII:**

# **PLANT AND ALGAE DERIVED COSMECEUTICALS AND NUTRACEUTICALS**

### **FLASH TALK**

#### **O45. Grape polyphenols may protect from diet induced obesity by preventing high-fat diet dampening of the hypothalamic-pituitary-adrenal axis**

Esther Mezhibovsky<sup>1,3</sup>, Kevin M. Tveter <sup>1</sup>, José A. Villa-Rodríguez <sup>1</sup>, Karen Bacalia<sup>1,3</sup>, Yue Wu<sup>1</sup>, Dushyant Kshatriya<sup>2,3</sup>, and Nicholas T Bello<sup>2</sup>, and Diana E. Roopchand <sup>1</sup>.

<sup>1</sup>Department of Food Science, Rutgers The State University of New Jersey, New Jersey Institute for Food Nutrition and Health and Rutgers Center for Lipid Research, New Brunswick, NJ 08901, USA. <sup>2</sup> Department of Animal Sciences, Rutgers The State University of New Jersey, New Brunswick, NJ 08901, USA. <sup>3</sup> Department of Nutritional Sciences, Rutgers The State University of New Jersey, New Brunswick, NJ 08901, USA. E-mail: em818@sebs.rutgers.edu

Chronic intake of a high-fat diet (HFD) induces weight gain due to impaired circadian hypothalamic-pituitary-adrenal (HPA) axis activity. Diet-induced HPA impairments are related to decreased levels of circulating corticosterone and altered glucocorticoid metabolism. Cholestasis has been shown to suppress HPA axis activity. HFD-fed mice supplemented with proanthocyanidin-rich grape polyphenols (GPs) have reduced body weight gain and hepatic steatosis, increased energy expenditure (EE), and altered bile acid (BA) gene regulation and profile. Here we investigate the potential of GPs to prevent HFD-induced disruptions to the HPA axis in association with BA changes. C57BL/6J male mice were fed HFD or HFD supplemented with 0.5% w/w GPs (HFD-GP) for 17 weeks. As previously seen, GP-supplemented mice had reduced weight gain, attenuated hepatic steatosis, and elevated EE. LC-MS analysis revealed a GP-induced reduction to circulating primary and secondary BAs, while qPCR showed increased hepatic expression of BA synthesizing enzymes, Cyp7a1 and Cyp27a1, and reduced ileal expression of BA transporter, Slc10a2, suggesting a mechanism for reduced cholestasis and BA leakage into circulation. Hormones involved in HPA axis activity, leptin and corticosterone, were increased in circulation of HFD-GP-fed mice while hepatic expression of 11 $\beta$ hsd1, an enzyme which converts inactive 11-dehydrocorticosterone to active corticosterone, was increased and corticosterone metabolizing enzymes, Srd5a1, and Akr1d1, were reduced. Expression of inflammatory markers Tnf $\alpha$ , Il6, and Il1b were reduced in both intestinal and hepatic tissue. We hypothesize that GPs improve metabolic health by protecting the liver from HFD-induced BA accumulation and the subsequent inflammation and disruptions to corticosterone and leptin regulation of EE. GP-supplementation may preserve diurnal regulation of the HPA axis and thus improve adaptations to excess dietary fat.

### **11<sup>th</sup> ANNUAL HYBRID CONFERENCE**

**American Council for Medicinally Active Plants (ACMAP)**

**EXPLORING THE UNEXPLORED: SCIENCE AND APPLICATIONS OF MEDICINAL PLANTS**

**JUNE 28 - JULY 2, 2022 | PUERTO RICO, USA**



## **SESSION VIII: PLANT AND ALGAE DERIVED COSMECEUTICALS AND NUTRACEUTICALS**

### **FLASH TALK**

#### **O46. Natural Products Research in Puerto Rico: Isolation of Bioactive Metabolites from Tropical Plants**

Claudia A. Ospina-Millán<sup>1</sup>. <sup>1</sup>Dept. of Natural Sciences and Mathematics, Inter American University of Puerto Rico, Bayamón, PR 00957. E-mail: [cospina@bayamon.inter.edu](mailto:cospina@bayamon.inter.edu)

According to the World Health Organization (WHO), ~80% of people in developing countries depend on traditional medicine for primary health care. Traditional herbal treatments have also yielded drug discovery efforts of purification and characterization of the active compound, as well as evaluation of safety, pharmacokinetics and efficacy of the drug candidate. In fact, 52% of the total molecules approved by FDA from 1981 to 2014, are natural products or their derivatives possessing structurally distinct families of chemical compounds. These data indicate that research in natural products continues to be a field of interest for the discovery of new drugs. However, less than 15% of plants have been assessed for bioactive potential, emphasizing a critical need for the discovery of novel bioactive compounds from unexplored plant species. Cancer is one of the major causes of death and research is focused on the discovery of novel drugs that are specific for tumor cells. It is expected that such drugs increase the effectiveness of treatment while reducing their side effects. Moreover, current therapeutic treatments of infectious diseases have limitations such as toxicity and low effectiveness. In addition, the emergence of strains resistance to current antifungal agents had led to the search for new drugs from natural sources. The overall goal of the proposed project is to discover and characterize novel compounds from higher plants with potential therapeutic application for cancer and anti-infective treatments. Our published and preliminary data showed that Caribbean plants of Croton, Guaiacum, Piper, Schinus and Simarouba genus provide an important source of new leads for in depth investigation as new anticancer and anti-infective therapeutics.



# **11<sup>th</sup> ANNUAL HYBRID CONFERENCE**


**American Council for Medicinally Active Plants (ACMAP)**

**EXPLORING THE UNEXPLORED: SCIENCE AND APPLICATIONS OF MEDICINAL PLANTS**

**JUNE 28 - JULY 2, 2022 | PUERTO RICO, USA**

## **POSTER SECTION**

### **VIRTUAL POSTER PRESENTATIONS**



**SCAN ME**

<https://bit.ly/Virtual-Poster-ACMAP2022>

**POSTERS RECORDED**

<https://bit.ly/posters-rec-2022>





## POSTER ABSTRACTS

### **P1. Isolation and characterization of novel plant-derived antimicrobial compounds from plantains.**

Christopher A. Sambolín-Pérez<sup>1</sup>, Juan A. Negrón-Berrios<sup>1</sup>, Rosalinda Aybar-Batista<sup>1</sup>; Michelle Cartagena-Rivera<sup>1</sup>; Alok Arun<sup>1</sup>. <sup>1</sup>Institute of Sustainable Biotechnology, Inter-American University of Puerto Rico, Barranquitas, PR 00794. E-mail: csam1223@intersg.edu; janegron@br.inter.edu

*Musa* spp. genus, specifically bananas and plantains have had a variety of applications throughout the history of mankind. This crop has been key to feeding population besides its variety of additional applications including medicinal properties. Currently, there is an emerging problem of bacterial resistant to commercially available antibiotics. Phytomedicine presents a sustainable solution to develop new compounds that can control bacterial growth. Pseudostem in banana reportedly contains an increased concentration of organic compounds, some of which have been partially characterized. This study aimed to study the antibacterial characteristics of extracts obtained from plantain pseudostem tissues. Organic extraction technique using the Soxhlet system was applied to produce organic and aqueous extracts. Further, these extracts were applied to *Pseudomonas aeruginosa* and *Escherichia coli* to analyze the bacteriostatic and bactericidal effect using Minimum Inhibitory Concentration (MIC) and Minimum Bactericidal Concentration (MBC) assays. Preliminary results indicate potential antimicrobial activity of plantain pseudostem extracts at 25.6 mg/ml on both tested bacteria. We are currently carrying out an in-depth validation of the antibacterial activity using a combination of microbial and biochemical assays. This study might help to identify and characterize novel sources of plant-derived antimicrobial compounds for a wider application in areas of human health.

### **P2. Genetic identification and biochemical characterization of *Marchantia* sp. From Puerto Rico**

Axel Alfredo Ferrer Vega<sup>1</sup>, Rosalinda Aybar Batista<sup>1</sup>, Ángel Núñez Marrero<sup>1</sup>, Juan Negrón Berríos<sup>1</sup>, Alok Arun<sup>1</sup>. <sup>1</sup>Institute of Sustainable Biotechnology, Inter American University of Puerto Rico, Barranquitas, PO Box 517, Puerto Rico, USA.  
E-mail: axelferrer89@gmail.com, alok\_arun@br.inter.edu

Bryophytes are considered the second largest group of land plants after angiosperms. These non-vascular plants play a prominent role in the maintenance of ecosystems as bioindicators to assess pollution in the environment and as medicinal plants to treat multiple disorders due to their many antioxidant properties. *Marchantia* is a bryophyte that has become a model plant for growing efficiently in vitro culture, the availability of the genome sequence and its dominance in the generation of haploid gametophytes make the plant suitable for genetic engineering. *Marchantia* from Puerto Rico was collected from the Carite State Forest, Toro Negro State Forest, and La Olimpia School Forest to identify its species, measure its antioxidant properties and determine the abundance of heavy metals. DNA barcoding method identified *Marchantia paleacea* as the species of interest from Puerto Rico. Measurement of antioxidant properties through assays such as flavonoid content, antioxidant capacity, and radical scavenging activity found that the sample from La Olimpia School Forest has the highest amount for a higher source of antioxidants than the others. The spectroscopy analytical technique determined a variation in the abundance of heavy metals in the samples of each forest that may represent contaminations in each sites. This study provides a pilot view of *M. paleacea* found in the forests of Puerto Rico, its antioxidant properties for a potential source of antioxidants and as a bioindicator for environmental pollution.



## POSTER ABSTRACTS

### P3. Bioprospecting of early-evolving land plants from Puerto Rico for pharmacological activities

Rosalinda Aybar Batista<sup>1</sup>, Gerardo Laureano<sup>1</sup>, Dem J. Santiago<sup>1</sup>, Juan Negrón Berríos<sup>1</sup>, Alok Arun<sup>1</sup>.

<sup>1</sup>Institute of Sustainable Biotechnology, Inter American University of Puerto Rico, Barranquitas, PO Box 517, Puerto Rico, USA. E-mail: raybar@br.inter.edu; alok\_arun@br.inter.edu

Plants are known for their unique biochemical properties. Recently, early evolving land plants are emerging model systems to extract metabolites. Bryophytes and pteridophytes are early evolving land plants that have been utilized in primitive eras for folklore medicines due to their medicinal properties. Research suggests that bryophytes and pteridophytes have been able to lower proliferation of cancer cells in addition to their antibiotic properties. Early evolving land plants from Puerto Rico remains to be explored. Our research aims to evaluate the medicinal properties of three bryophyte species and two pteridophyte species, by estimating the number of flavonoids, total polyphenol, and antioxidant capacity. Our preliminary results show that the tissues from pteridophytes have more amount of total flavonoids and total polyphenol content. Samples from pteridophytes showed 214.44µg/mL of total flavonoid content as compared to 71.09µg/mL in bryophytes. Similarly, pteridophytes show an average of 171.25µg/mL of total polyphenol content in comparison with Bryophytes that showed an average of 62.23µg/mL. Furthermore, we determined the antimicrobial effect of these plants on four different gram-positive and gram-negative bacterial strains, using the methanolic extracts of each plant obtained by Soxhlet extraction method. This assay demonstrated once again that pteridophytes have effect inhibiting the growth of some of these bacteria compared to the bryophytes in study. Our study provides a preliminary data to carry out an in-depth bioprospecting of early evolving land plants from Puerto Rico.

---

### P4. Cytotoxicity effect of medicinal plants used by the Moroccan population against Covid-19

Houda ZAHER<sup>1,2</sup>, Latifa TAJOUNTE<sup>2</sup>, Sanaa LEMRABET<sup>2</sup>, Noureddine HAMAMOUCHE<sup>3</sup>, Hicham OUMZIL<sup>2</sup>, Bouchaib BENCHARKI<sup>1</sup>. <sup>1</sup>Laboratory of Agri-food and Health, Faculty of Science and Technology, University Hassan 1st, Settat, Morocco. <sup>2</sup>Department of Virology, National Institute of Hygiene, Ministry of Health, Rabat, Morocco. <sup>3</sup>Laboratory of Plant Biotechnology and Physiology, Faculty of Sciences, University Mohammed V, Rabat, Morocco. E-mail: h.zaher@uhp.ac.ma

The present study was conducted to test the cytotoxicity effect of essential oils from three different medicinal plants used by the Moroccan population against Covid-19; *Melissa officinalis*, *Zingiber officinalis* and *Eucalyptus globulus*. Plant essential oils were obtained from plant leaves by hydrodistillation and their cytotoxicity was determined using VERO-E6 cells (ECACC-code 84121903) in a standard red dye test. These plants were selected on the basis of an online ethnobotanical survey on medicinal plants used by people in Morocco against Covid-19. The results of the neutral red assay on VERO-E6 cells showed that essential oil from *M. officinalis*, *Z. officinalis* and *E. globulus* exhibit moderate cytotoxic activities with a cell viability of 50%. The extracts from these plants are currently being tested against Covid-19. Cytotoxicity studies of medicinal plants used against Covid-19 is The results provide an attractive perspective for further exploration of their antiviral and anti-inflammatory potential to combat viral respiratory infections.



## POSTER ABSTRACTS

### **P5. Study of the chemical composition of the hexane extract from Simarouba**

Neymar Maldonado-Vargas<sup>1</sup>, Paola C. Nieves-Oquendo<sup>1</sup> and Claudia A. Ospina-Millán<sup>1</sup>. <sup>1</sup>Dept. of Natural Sciences and Mathematics, Inter American University of Puerto Rico, Bayamón, PR 00957. E-mail: nmaldonado6965@interbayamon.edu; cospina@bayamon.inter.edu

The Simaroubaceae family comprise around 32 genera and more than 170 species of trees and brushes distributed throughout the tropical regions of America, Asia, Africa, and Australia. This family is characterized by the presence of quassinoids that are responsible for antimalarial, antiviral, insecticide, and antitumor activities. In our ongoing efforts to discover new compounds with anticancer activity, we tested the hypothesis that leaf extracts from Simarouba showed the highest antiproliferative effects against breast cancer cell lines (Claudio et al., 2015). From the chloroform extract, we reported the isolation and cytotoxic activities of the quassinoid Simalikalactone D (SKD) which showed potent in vitro cytotoxicity with an IC50s of 58 to 65 nM on ovarian and breast cancer cell lines (Mendez et al., 2020). The hexane extract showed an inhibition potential at a concentration of 24 ng/mL against breast cancer cells. However, the chemical composition is unknown. The goal of our work is to isolate the secondary metabolites with less polarity and evaluate their anticancer activity. The <sup>1</sup>H-NMR spectrum of the extract showed the presence of aliphatic, vinylic,  $\alpha$ -heteroatoms and vinylic protons. This crude extract was chromatographed on Silica gel with a mixture of 10% of ethyl acetate in hexane to obtain 20 fractions. The fractions are being analyzed by TLC and NMR to identify the chemical components.

### **P6. Volatile extracts from blackberries, black raspberries, and blueberries inhibits proliferation of human lung cancer A549 cells through apoptosis**

Inah Gu, Cindi Brownmiller, Luke Howard, Sun-Ok Lee\*. Department of Food Science, Division of Agriculture, University of Arkansas, Fayetteville, AR 72704, USA. E-mail: inahgu@uark.edu; cbrownm@uark.edu; lukeh@uark.edu; sunok@uark.edu

Berries are one of the most common fruits in our diet and are rich in vitamins, minerals, dietary fibers, and especially polyphenols and volatiles. Berry volatiles are low molecular weight compounds that are in charge of the berry aroma and flavor. There have been many studies conducted to identify the chemical composition of berry volatiles for developing consumer acceptability. However, there are lack of information available on the health-promoting properties of berry volatiles. In this study, we investigated the apoptotic effect of blackberry, black raspberry, and blueberry volatile extracts on human lung adenocarcinoma A549 cells. Cells were treated with three dilutions (2-, 4-, and 8-fold diluted) of three berry volatile extracts (BVEs) for 12, 24, and 48 h. The proliferation of A549 was measured by the MTS assay. Changes in the cell cycle and apoptosis were examined by using cell sorting analysis, flow cytometry, and cell death detection ELISA kit. The apoptotic mechanism was further investigated by western blot analysis. Statistical analysis was performed by using one-way ANOVA, followed by Tukey's multiple comparisons to determine significant differences at  $p < 0.05$ . All two-fold diluted BVEs significantly inhibited the proliferation of A549 after 48 h by inducing apoptosis after 24 h ( $p < 0.05$ ). Two-fold diluted BVEs treatment for 48 h also increased the cell population in G0/G1 phase (64.9-73.1%) compared to control (60.3%). Flow cytometric quantification of apoptosis identified the most early apoptotic cells in blackberry volatile extracts. Similarly, blackberry volatile extracts significantly inhibited the level of poly ADP-ribose polymerase (PARP), procaspase-9, and procaspase-3 compared to control ( $p < 0.05$ ). These results showed that berry volatile extracts from three berries have the anti-proliferative effect on human lung adenocarcinoma cells via apoptosis, suggesting that berry volatiles from blackberry, black raspberry, and blueberry may have potential anticancer activity through the apoptosis in lung cancer.



## POSTER ABSTRACTS

### **P7. Gene expression changes elicited by *Nigella sativa* compounds in glioblastoma cells**

Rachana Pandey<sup>1</sup>, Purushothaman Natarajan<sup>1</sup>, Umesh Reddy<sup>1</sup>, Wei Du<sup>2,3</sup>, Cristian Sirbu<sup>2,3</sup>, Moussa Sissoko<sup>4</sup>, Gerald R. Hankins<sup>1</sup>. <sup>1</sup>West Virginia State University Department of Biology and Biotechnology Graduate Program, Institute WV 25112, <sup>2</sup>Charleston Area Medical Center Cancer Center,<sup>3</sup> Charleston Area Medical Center Institute for Academic Medicine, Charleston, WV 25302. <sup>4</sup>Katmay Oncology Group, Anchorage, Alaska, 99508. E-mail: rpandey1@wvstateu.edu; ghankins@wvstateu.edu

Approximately 1,700,000 new cases of cancer are diagnosed in the United States, and around 600,000 will die of the disease. While cancer survival rates have improved over time, the survival rates are still low for many cancers. Examples include the deadly brain cancer glioblastoma, for which there is less than a 10% five-year survival post-diagnosis and pancreatic cancer with a five-year survival rate of only 6%. The seeds of *Nigella sativa* (family Renuculaceae) have been used for culinary and multiple medical purposes for over 2000 years. These are commonly known as “black seed” or “black cumin”. In modern times pharmacologically active compounds have been identified in the oil from the seed. The most abundant and most well studied is the monoterpene, thymoquinone. Although several mechanisms for the anti-cancer activity of thymoquinone have been identified, we present here the first RNA sequencing study directed towards comprehensively identifying pathways affected by thymoquinone in these deadly brain cancers. In this case, we examine the effects of gene expression on A172 human glioblastoma cells. We have performed deep RNA Sequencing and generated 25-30 million reads per sample using next-generation sequencing. These reads are being mapped to the human reference genome to identify genes that are differentially expressed between untreated and thymoquinone treated cells. We will present differentially expressed genes and pathways that may be relevant to the response of the glioblastoma cells to *Nigella sativa* compounds.

---

### **P8. Arachidin-1 induces apoptosis and growth inhibition in triple-negative breast cancer cells**

Alexx Weaver<sup>1</sup>, Sepideh Mohammadhosseinpour<sup>1</sup>, Fabricio Medina- Bolivar<sup>1,2</sup>. <sup>1</sup>Arkansas Biosciences<sup>2</sup>Institute Dept. of Biological Sciences, Arkansas State University, Jonesboro, AR 72401. E-mail: alexx.weaver@smail.astate.edu; fmedinabolivar@astate.edu

Triple-negative breast cancer (TNBC) is an aggressive, fast-spreading, and hormonal treatment-resistant form of breast cancer. These factors lead to worse prognoses for TNBC patients, stressing the need for investigating alternative treatment options for this type of cancer. Plant natural products, such as the prenylated stilbenoid arachidin-1 (A-1) from peanut, have shown cytotoxicity to certain cancer cells while showing no significant cytotoxicity to non-cancerous cell lines. The goal of this study was to examine the apoptotic effects of A-1 alone and in combination with paclitaxel, a chemotherapeutic drug that causes adverse side effects. The apoptotic effects of these treatments were studied in MDA-MB-436 and MDA-MB-231 via western blotting through protein expression levels of PARP, caspase-8, caspase-9, and survivin. Vinculin and GAPDH were protein-loading controls. Flow cytometry was used to detect apoptosis and cell cycle arrest in the treated cells.

Results showed that A-1 decreased the IC<sub>50</sub> of paclitaxel in TNBC cells, and consequently increased the expression of cleaved PARP and cleaved caspase-9. The amount of survivin, an apoptosis inhibitor protein, decreased at increased A-1 concentrations. In addition, flow cytometry analysis showed that A-1 in combination with paclitaxel led to cell cycle arrest in G2-M stage and induction of apoptosis in the treated cells. These results support an increase of apoptotic effects of paclitaxel when combined with A-1. This accentuates the importance of expanding our understanding of the connection between stilbenoids and triple-negative breast cancer cells. In future research, we will study the effects of A-1 on signaling pathways in TNBC cell lines.



## POSTER ABSTRACTS

### P9. Antimalarial activity of Puerto Rican tropical plants

Angenil Garcia-Ortiz<sup>1</sup>, Emilee E. Colón-Lorenzo<sup>2</sup>, Adelfa E. Serrano-Brizuela<sup>2</sup> and Claudia A. Ospina-Millán<sup>1</sup>. <sup>1</sup>Dept. of Natural Sciences and Mathematics, Inter American University of Puerto Rico, Bayamón, PR 00957 and <sup>2</sup>Dept. of Microbiology and Medical Zoology, School of Medicine, University of Puerto Rico, Medical Sciences Campus, PR 00936. E-mail: agarcia1903@interbayamon.edu; cospina@bayamon.inter.edu

Malaria, as a major global health problem, continues to affect many people in developing countries worldwide. There is a critical need to discover compounds from new sources to overcome the problem of drug-resistant Plasmodium strains. Plants continue to be one of the most important sources in the search of new antimalarials. This study aimed to investigate the antimalarial activity of the crude extract, solvent fractions, pure compounds, and tea preparations from Puerto Rican tropical plants. In a preliminary study, one of the plants specie tested positive to antiplasmodial activity against *P. berghei*. The crude extract of this plant was chromatographed in silica gel with a mixture of 2% of methanol in chloroform to obtain 8 fractions. Fractions 4 and 6 were purified by reversed phase chromatography with a mixture of 45% of water in methanol to obtain compounds A and B. Compounds A and B showed a potent antiplasmodial activity at EC<sub>50</sub> of 15 and 18 nM against *P. berghei*, respectively. This study suggests that tropical plants from Puerto Rico are promising sources of compounds that might be further investigated as novel drugs against malaria.

---

### P10. Antioxidant and antibacterial acclivities of *Dittrichia viscosa* extracts

Bentarhlia Noura<sup>1,2</sup>, El Monfalouti Hanae<sup>1</sup>, Taghzouti Khalid<sup>2</sup>, Nouredine HAMAMOUC<sup>3</sup>, Abboussi Oualid<sup>2</sup>, Kartah Badreddine<sup>1</sup>. <sup>1</sup>Laboratory of Plant chemistry, organic and bioorganic synthesis, Faculty of Sciences, Mohammed V University in Rabat, Morocco. <sup>2</sup>Physiology and Physiopathology Team, Faculty of Sciences, Genomic of Human Pathologies Research Centre, Mohammed V University in Rabat, Morocco. <sup>3</sup>Laboratory of Plant Biotechnology and Physiology, Faculty of Sciences, University Mohammed V, Rabat, Morocco. E-mail: noura.bentarhlia@um5r.ac.ma

*Dittrichia viscosa* is a medicinal plant traditionally and widely used to treat several diseases such as diabetes fever and inflammation. In morocco its fresh leaves are crushed and given for topical application. The present study aim to evaluate the antimicrobial and antioxidant potentials of ethanol, aqueous, hexane, ethyl acetate and n-butanol extracts of the aerial part of *Dittrichia viscosa*. Antimicrobial activity was investigated against Gram-positive bacteria (*Bacillus subtilis* and *Staphylococcus aureus*), Gram-negative bacteria (*E. coli* and *Pseudomonas aeruginosa*) and two fungi (*candida albicans*, *saccharomyces cerevisiae*) using agar disc diffusion and dilution methods. Antioxidant activity was assessed by 1, 1-diphenyl-2-picrylhydrazyl (DPPH) free radical scavenging and total antioxidant capacity (TAC). The result showed a significant Scavenging activity against DPPH free radicals, total antioxidant capacity (TAC) as well as antimicrobial activity at concentration of 5 mg/ml. The results have proven the potential of antimicrobial and antioxidant natural alternative to preserve foodstuff avoiding healthy hazards of chemical antimicrobial agent applications and natural antioxidant source.



## POSTER ABSTRACTS

### P11. Study of the chemical composition and biological activities of *Piper* sp.

Natalia Rivera-Colón<sup>1</sup>, Pablo E. Vivas-Mejía<sup>2</sup>, Lilliam Villanueva-Alicea<sup>2</sup>, José Rodríguez-Medina<sup>2</sup> and Claudia A. Ospina-Millán<sup>1</sup>. <sup>1</sup>Dept. of Natural Sciences and Mathematics, Inter American University of Puerto Rico, Bayamón, PR 00957 and <sup>2</sup>Dept. of Biochemistry, University of Puerto Rico, Medical Sciences Campus, PR 00936. E-mail: nrivera2619@interbayamon.edu; cospina@bayamon.inter.edu

*Piper* genus are aromatic plants belonging to Piperaceae family. These plants are native of Puerto Rico and the Virgin Islands. It has been found that compounds isolated from different *Piper* specie have anticancer and antifungal activities. The objective of this work is to evaluate the anticancer and antifungal properties of *Piper* genus and isolate the chemical compounds responsible for biological activity. The plants were collected, dried, and extracted with a mixture of CH<sub>2</sub>Cl<sub>2</sub>-MeOH (1:1). The resulting crude extract was suspended in water and extracted with solvents of different polarities. The crude extract was analyzed by nuclear magnetic resonance (NMR) and tested for the anticancer and antifungal activities. Our preliminary results showed that this extract has anticancer antiproliferative activity against the A2780CP20 ovarian and MCF-7 breast cancer cell line cancer cell lines at a concentration of 0.18 and 0.14 mg/mL, respectively. In addition, the extracts were active against *Saccharomyces cerevisiae* at a concentration of 500 µg/mL. The hexane extract was chromatographed over silica gel with a mixture of 15% of ethyl acetate in hexane to obtain 16 fractions. The NMR analysis of fractions 11 to 13 showed the presence of signals corresponding to aromatic compounds. We are working with the characterization of the chemical compounds.

---

### P12. Chemical Analysis and Biological Properties of the Native Plant *Schinus terebinthifolius*

Sabrina Elías-Rodríguez<sup>1</sup>, Angenil García-Ortiz<sup>1</sup>, Lilliam Villanueva-Alicea<sup>2</sup>, José Rodríguez-Medina<sup>2</sup> and Claudia A. Ospina-Millán<sup>1</sup>. <sup>1</sup>Dept. of Natural Sciences and Mathematics, Inter American University of Puerto Rico, Bayamón, PR 00957 and <sup>2</sup>Dept. of Biochemistry, University of Puerto Rico, Medical Sciences Campus, PR 00936. E-mail: selias3731@interbayamon.edu; cospina@bayamon.inter.edu

Fungal diseases represent a substantial cause of infections in plant and animal species as well as a public health problem. In addition, the emergence of strains resistant to current antifungal agents has led to search for new drugs from natural sources. *Schinus terebinthifolius* is a plant belonging to the Anacardiaceae family native to South America, which is also found in tropical regions of the United States and the Caribbean. In traditional medicine, it has been used as an antimicrobial and anti-inflammatory agent. It is known that the chemical composition of a plant varies according to where it is collected. To our knowledge, the *Schinus* sp. that grow in Puerto Rico have not been studied. The crude extract from the leaves was tested for determination of antifungal activity on wild-type *Saccharomyces cerevisiae* yeast strains using a viability assay on yeast agar plate cultures. Preliminary results showed that this extract presented 100% antifungal activity at a concentration of 500 µg/mL in the first 72 hours. Based on this observation, our objective is to assess the antifungal potential of *Schinus* extracts and the purified compounds. For this purpose, the plant was collected, dried, and extracted with a mixture of CH<sub>2</sub>Cl<sub>2</sub>-MeOH (1:1). The resulting crude extract was suspended in water and extracted with solvents of different polarities. Fraction 15 of the hexane extract was chromatographed on silica gel with a mixture of 10% of ethyl acetate in hexane to obtain 17 fractions. The NMR analysis of fractions E and F revealed the presence of aliphatic, allylic, α-heteroatoms and vinylic protons. We are working with the characterization of the chemical compounds.

## POSTER ABSTRACTS

### **P13. Production of cajaninstilbene acid in hairy root cultures of pigeon pea (*Cajanus cajan*) using an optimized elicitation procedure**

Gaurav Gajurel<sup>1,2</sup>, Luis Nopo-Olazabal<sup>1</sup>, Emily Hendrix<sup>1</sup>, Fabricio Medina-Bolivar<sup>1,3</sup>. <sup>1</sup>Arkansas Bioscience Institute, <sup>2</sup>Molecular Biosciences Graduate Program, and <sup>3</sup>Department of Biological Sciences, Arkansas State University, Jonesboro, AR 72401.

E-mail: gaurav.gajurel@astate.edu; fmedinabolivar@astate.edu

Cajaninstilbene acid (CSA) is a prenylated stilbenoid derivative that exhibits anti-inflammatory, anti-bacterial, and neuroprotective properties. Hairy root cultures of pigeon pea (*Cajanus cajan*) were developed via *Agrobacterium rhizogenes*-mediated transformation to establish a production platform for this bioactive compound. The cultures were co-treated with methyl jasmonate, cyclodextrin, hydrogen peroxide, and magnesium chloride to enhance the production of CSA. The majority of CSA accumulated in the culture medium. CSA was purified from extracts of the culture medium by semi-preparative HPLC, and its identity was confirmed by tandem mass spectrometry. After 144 h of elicitation in 12-day old hairy root cultures, the total yield of CSA was  $8058.61 \pm 445.78$   $\mu\text{g/g}$  DW. The yield of CSA in the elicited hairy root culture was approximately 277-fold higher than in the non-elicited hairy root culture. The difference in phenotype of elicited versus non-elicited pigeon pea hairy roots was studied using scanning electron microscopy. The non-elicited hairy roots had uniform surfaces whereas the elicited hairy roots had non-uniform shapes. Pigeon pea hairy roots provide a sustainable platform for producing CSA.

### **P14. Phytochemicals from PhytoMed Honey with high docking affinity with SARS CoV-2 proteins and the traditional medicinal uses of their plant sources**

VValma Jessamy<sup>1</sup> and Biswaranjan Pradhan<sup>2</sup>. <sup>1</sup>Caribbean Institute for Sustainable Island Living, St. George's, Grenada. <sup>2</sup>Indian Institute for Technology, India.

E-mail: vjessamy@jecocaribbean.com

PhytoMed Honey® is a varietal of honey which was naturally produced by developing a proprietary hive management process and proprietary planting scheme of tropical medicinal plants native to the Caribbean as the nectar and pollen source for honeybees *Apis Mellifera*. PhytoMed Honey was administered to PCR positive Covid-19 patients in a Human Covid Trial and patients symptoms improved after 72 hours with second PCR test negative.

Using HRMS 176 phytochemicals were identified in the honey and 35 are known to have anti-viral activity against other viruses. Molecular docking studies of the antiviral phytochemicals in PhytoMed Honey revealed the mechanism by which the honey was able to express activity against SARS-CoV-2. Twenty-three phytochemicals with the highest docking affinity ( $>-7\text{kal/mol}$ ) with main protease (7alh) were linked to their plant source. Pharmacological studies have validated the use of these plants in traditional herbal medicine for treatment of a range of disease conditions. The combination of phytochemicals in PhytoMed Honey present a novel formulation for use of the medicinal plants.



## POSTER ABSTRACTS

### **P15. Chemical Diversity Assessment and Ethnobotanical Survey of Important Indian Medicinal Plants**

K M Patel<sup>1</sup> and Satyanshu Kumar<sup>2</sup> <sup>1</sup>Principal (Retd.) Municipal Arts and Urban Bank Science College, Mehsana-384002 Gujarat, India and <sup>2</sup>ICAR-Directorate of Medicinal and Aromatic Plants Research, Anand-387310, Gujarat, India. E-mail: kmpatel.flyfortune@gmail.com

India is one of the world's 12 biodiversity centres with the presence of over 45000 different plant species. Out of these, about 15000-20000 plants have good medicinal value. About 7000-7500 species are used for their medicinal values by traditional communities. Plant secondary metabolites belong to group of naturally occurring compounds and its content in plant is subjected to environmental influences. By using scientific evaluation of chemical diversity of plants, their medicinal properties could be established. Ethnobotanical study establishes the relationship between humans and plants. Ethnobotany has played important role in the development of new drugs for centuries. However, even today, ethnobotanical knowledge is still transmitted from generation to generation chiefly by mouth. The rising global demand for plant based medicines also presents existential threats to many plant species. Therefore, ethnobotany may have important role in sustainable supply of some of the important medicinal plants. In addition to that it can also be of use to find a new medicinal recipe which could be used to treat diseases for which no standard therapy have been reported. Bioactive compounds of plant origin are those secondary metabolites possessing desired health/ wellness in man and animals. Secondary metabolites are chemically and taxonomically extremely diverse compounds with obscure function. In the present study, nine high value endangered medicinal plants of India namely *Aconitum heterophyllum* (Atish), *Ephedera foliata* (Ephedera), *Malaxis acuminata* (Jeevak), *Pterocarpus marsupium* (Vijaysar), *Podophyllum hexandrum* (Himalayan Mayapple), *Rauvolfia serpentina* (Sarpagandha), *Rheum emodi* (Rewand chini), *Swertia chirata* (Chirayata), *Valeriana jatamansi* (Jatamansi) have been selected for the assessment chemical diversity in terms of the major phytochemical. These species are not in cultivation and collected from nature, therefore, ecological studies needs to be brought to the forefront in order to protect their in-situ conservation. Outcome of the present study may be useful to identify the active constituents to discover new chemotypes, as promising sources of drugs. Also usefulness of ethnobotany research lies in the increased consumer preference to traditional medicine and also increased demand for natural medicines.

---

### **P16. Some important medicinal orchids plants with their medicinal uses from Northwest Himalayan (Uttarakhand)**

Dr.Uma Rani. S.S.D.P.C .Girls (P.G.) College, Roorkee, Uttarakhand, India.  
E-mail: drumarani10@gmail.com

The Himalayan range has a rich heritage of knowledge on orchids' plant-based therapy. Medicinal orchid plants play a major role in livelihood from all over the world. Uttarakhand, a Himalayan state of India also depends on medicinal orchid plants for medicine and traditional therapy. Orchids are nature's most extravagant group of flowering plants distributed throughout the world from tropics to high alpine. They exhibit an incredible range of diversity in the size, shape, and color of their flowers. Though orchids are grown primarily as ornamentals, many are used as herbal medicines, food, and others have cultural value by different cultures and tribes in different parts of the world. Orchids have been used in many parts of the world in traditional healing systems as well as in the treatment of a number of diseases since ancient times. Though Orchidaceae is regarded as the largest family of the plant kingdom, few studies have been done regarding their medicinal properties. Linking the indigenous knowledge of medicinal orchids to modern research activities provides a new reliable approach, for the discovery of novel drugs much more effectively than with random collection. Many of these orchids face the extreme danger of extinction due to over-exploitation and habitat loss. Plant tissue culture could be one of the most suitable alternative tools to minimize the pressure on the natural population of medicinal orchids and their sustainable utilization. In this scenario, we try to document some orchid medicinal plants with their medicinal properties from this state.

## POSTER ABSTRACTS

### **P17. The Evaluation of Atmospheric Pressure Plasma, Hot Water, and Heat Pads on the Sprouting, Yield and Curcumin Content of Turmeric (*Curcuma longa*).**

Trang Pham<sup>1</sup>, Khadejah Scott<sup>1</sup>, Lam Duong<sup>1</sup>, Suresh Kumar<sup>1</sup>, DeAnthony Price<sup>1</sup>, Mei Wang<sup>2</sup>, Charles Cantrell<sup>2</sup>, and Srinivasa Rao Mentreddy<sup>1</sup>.<sup>1</sup>Department of Biological and Environmental Sciences, Alabama A&M University, Huntsville, AL 35811; <sup>2</sup>USDA-ARS-NPURI, University, MS. E-mail: [srinivasa.mentreddy@aamu.edu](mailto:srinivasa.mentreddy@aamu.edu)

Curcuminoids produced in the rhizomes of *Curcuma* spp., have antioxidant and anti-inflammatory activity. The U.S. imports 90% (\$51.7 million) of turmeric from Asian Countries, mainly India. The U.S. herbal manufacturers seek domestically produced turmeric to replace imports. The study's objective was to determine the effects of hot water (HW) and heat pads (HP) at three temperature levels, and exposure to atmospheric pressure plasma (APP) for 90 seconds on days to sprouting, growth, yield, and curcumin levels of turmeric. The rhizomes of three turmeric varieties, CL11, VN39, and VN50, subjected to i) HW at room temperature (HWCo), 35°C (HW35), 45°C (HW45), and 55°C (HW55) for 30 min; ii) HP at 24°C (HP24), 29°C (HP29), and 35°C (HP35); and iii) exposure to APP for 90 seconds, and untreated (HPCo) were grown in the greenhouse. The plants were then transplanted on to field plots. Curcuminoids, curcumin, demethoxy-curcumin, bisdemethoxy-curcumin, and aR-Turmerone were determined using HPLC. The HP treatment promoted early sprouting among the three varieties, with 30% of the rhizomes sprouting 4 to 6 days earlier than the rhizomes in other treatments. The APP treatment enabled 100%, 85%, and 80% of the rhizomes of CL11, VN39, and VN50, respectively, to sprout within 65 days after treatment. The plants in HP were taller than those in other treatments. HW35CL11 produced the highest yield (11.5 ton/ha) and HW45VN50 had the highest curcuminoid content (6.42%). In all varieties, treatments increased curcuminoid levels (4 to 7%) compared to Control (3.8 to 6%). The HP treatment promoted early sprouting whereas HW treatment enabled higher yields. Soaking rhizomes in water for 30 min at room temperature improved yield and curcumin content compared to control no treatment. Therefore, using HP or HW35, HW45 for 30 minutes offers the potential for extending the crop growing season, high yield and high curcuminoid content of turmeric.

### **P18. Turmeric (*Curcuma longa* L.): An Important Medicinally Active Plant Investigated for Ease of Germination Using a Laboratory Method**

ASparks<sup>1</sup>, D. Roberson, A. and B. K. Biswas,<sup>1</sup>MS Biotechnology Research Scholar, College of Agriculture, Family Sciences and Technology, Fort Valley State University, Fort Valley, GA 31030. E-mail: [biswasb@fvsu.edu](mailto:biswasb@fvsu.edu)

Turmeric (*Curcuma* spp.), native to southeast Asia, has been used in Indian culture for more than 4,000 years for multiple uses, like medicinal, culinary, cosmetics, and as a dye. Research worldwide has shown that curcumin, an active ingredient of turmeric, has potential to cure cancer, arthritis, inflammation, Alzheimer and so on. Therefore, to enable turmeric production in the USA and to benefit US consumers and growers, this experiment was carried out to examine the germination potential of turmeric in the laboratory. In this study, two different methods were used: i) in Ziplock bags and ii) in a rocker shaker box. Turmeric rhizomes were collected from various sources: 1. hydroponics grown germplasm, 2. raised beds, and 3. field plots. The rhizomes were placed in the rocker shaker box and in the Ziplock bags filled with 50-100 mL of distilled water, so that rhizomes were soaked well, but not completely submerged in water. In both experiments, rhizomes sprouted within 2-3 days and grew to 2.5 cm within 7-9 days. The rhizomes in the rocker shaker boxes sprouted within 2 days and grew faster than the rhizomes in the plastic bags. The Ziplock bag turmeric sprouted a day later- on day 3 and grew slower than those in the rocker shaker box. The experiment will be repeated and the data will be present at the ACMAP conference.



## POSTER ABSTRACTS

### **P19. Clonal propagation of selected lines of stevia (*Stevia rebaudiana* Bertoni) via in vitro micropropagation**

Maria de Lourdes Tapia y Figueroa and Luz Raida Gómez Pando. Departamento de Fitotecnia, Facultad de Agronomía, Universidad Nacional Agraria La Molina, Peru  
E-mail: ltapia@lamolina.edu.pe

*Stevia* (*Stevia rebaudiana* Bertoni) is a species native to Paraguay and Brazil that is used as a sweetener or natural sweetener. It also has beneficial effects on health for its medicinal properties that contribute to reducing problems of glycemia and hypertension. Vegetative propagation is possible by cuttings but it has the problem of spreading plants contaminated with diseases. Therefore not enough quality plants are produced for large-scale production. Furthermore, the process is labor-intensive and the seeds produced by sexual propagation have low germination. One way to overcome these limitations is to propagate stevia by in vitro micropropagation, and the present work resulted in micropropagation and evaluation of three selected lines IBT (IBT1, IBT2 and IBT3). Among these, line IBT1 presents the highest values for seedling height, number of leaves and fresh weight and the IBT2 line has the lowest values. In the acclimatization stage, IBT1 line had a survival of 85% whereas the IBT2 line 80% and the IBT3 line showed 89%.

### **P20. Antiarthritic potential of crude extract of *Firmiana colorata* (roxb.) flowers in FCA induced arthritis in rats**

K. M. Patel <sup>1</sup>, J. O. Bhangale <sup>2</sup>, P. J. Bhangale <sup>2</sup> <sup>1</sup>Principal (Retd.), Municipal Arts and Urban Science College, Mehsana 384002, Gujarat, India. <sup>2</sup>Smt. N. M. Padalia Pharmacy College, Navapura, Ahmedabad, 382 210 Gujarat, India. E-mail: kmpatel.flyfortune@gmail.com

The present work deals with antiarthritic potential of crude extract of *Firmiana colorata* (ROXB.) flowers on FCA induced arthritis in rats. Different extracts from petroleum ether (PEFC), ethyl acetate (EAFC) and alcohol (AFC) were administered at a dose 100, 200 and 400 mg/kg, orally, from day 13 to day 21. Symptoms of Arthritis was assessed by change in body weight, change in arthritic score, change in paw volume and change in ankle diameter on day 0, 3, 6, 9, 12, 15, 18 and 21. At the end of study, animals were anesthetised and blood was collected for the estimation of various hematological parameters such as hemoglobin content, total RBA, WBC, ESR count, CRP level and TNF alpha level. There was a significant increase in body weight, reduction in arthritic score, paw volume and ankle diameter in PEFC (200 and 400 mg/kg); EAFC and AFC (400 mg/kg, p.o.) treated animals. Above finding was confirmed by hematological results as, significant improvement in the levels of Hb and RBC count, and suppressed WBC count, ESR count, CRP and TNF alpha levels found in PEFC (200 and 400 mg/kg); EAFC and AFC (400 mg/kg, p.o.) administered arthritic group. Our findings showed a significant antiarthritic activity of *Firmiana colorata* (ROXB.) flowers against FCA induced arthritis in rats.



## POSTER ABSTRACTS

### **P21. Using Gold Nanoparticles for the Sensitive Detection of Aflatoxin B1 in Peanuts**

Amber Davenport<sup>1</sup> and Hari Singh<sup>1</sup>. <sup>1</sup>Nanotechnology Laboratory, College of Agriculture, Family Sciences and Technology, Fort Valley State University, Fort Valley, GA 31030.

E-mail: [adavenp6@wildcat.fvsu.edu](mailto:adavenp6@wildcat.fvsu.edu); [singhh@fvsu.edu](mailto:singhh@fvsu.edu)

It is estimated that more than 25% of the world's food supply is contaminated by mycotoxins and up to 4.5 billion people are exposed annually. Mycotoxins, or toxins produced by fungi, are naturally occurring contaminants produced by the *Aspergillus*, *Penicillium* and *Fusarium* fungi and are known to contaminate food supplies and animal feeds throughout the world. Aflatoxins are highly toxic mycotoxins and are found on agricultural crops such as maize (corn), peanuts, cottonseed, and tree nuts and depending on the level of exposure, it can result in growth retardation, immune-suppression, liver damage or at worst, death. Peanuts are a major source of food throughout the world, consumed either as peanut butter, crushed and used for oil, or simply consumed as a snack. The total U.S. peanut production in 2021 measured more than 6.3 billion pounds of peanuts and more than half of them were grown in Georgia. In terms of income, aflatoxins are a major limitation to accessing lucrative export markets leading to loss of income for farmers. There are at least 20 known aflatoxins and of the aflatoxins, aflatoxin B1 (AFB1) is the most toxic and the most potent naturally occurring chemical liver carcinogen. Most human exposure comes from contaminated nuts, grains, and their derived products. The toxins can be present even if there is no visible mold on the food, which is why early detection is key. There have been methods used to detect mycotoxins, in the past, but they require the use of expensive, large machines or have a low sensitivity. The aim of this research is to develop a nanobiosensor for the rapid, sensitive, and selective detection of aflatoxin B1 (AFB1) in peanuts using magnetic beads, carbon probes, and gold nanoparticles.

### **P22. Use of *Aspergillus oryzae* for the enrichment of zinc and selenium by using solid-state fermentation**

Rashmi Rangaswamaiah and Sun-Ok Lee. Department of Food Science, Division of Agriculture, University of Arkansas, Fayetteville, AR 72704, USA. E-mail: [rrangasw@uark.edu](mailto:rrangasw@uark.edu); [sunok@uark.edu](mailto:sunok@uark.edu)

Zinc and selenium are essential minerals for human health. Studies suggest that they enhance immunity, antioxidant effect, growth, reproductive performance, and the ability to resist disease. They are naturally found in many food sources such as dairy, poultry, cereals, and other grain products. However, the risk of zinc and selenium deficiency has been recognized in a few countries. This could be due to the lack of zinc and selenium content in the crops' soil. Meanwhile, antinutrients in the food source also inhibit proper absorption of minerals into the human body. But these antinutrients can be reduced by the fermentation process. The fermentation is carried out by yeast, bacteria, fungi, or a combination of these. In this study, solid-state fermentation (SSF) was applied to determine the amount of zinc and selenium uptake by *A. oryzae*. The sorghum grain (substrate) was treated with *A. oryzae* and supplemented with the different concentrations of zinc or selenium. After 7 days of fermentation, the samples were analyzed with Inductively Coupled Plasma Mass Spectrometry (ICP-MS). The results showed an exponential improvement in zinc content in the fermented samples, from the initial value of 4.5 mg/g to 158 mg/g, and selenium content increased from 0.19 mg/g to 0.8 mg/g. Data indicate that zinc and selenium uptake by *A. oryzae* from substrate significantly ( $P < 0.05$ ) varied by added concentration. Overall, this research shows that *A. oryzae* can uptake and increase the amount of zinc and selenium through SSF and the resulting product may use as zinc or selenium-rich ingredient for functional food products.



## POSTER ABSTRACTS

### **P23. Artificial neural network models for the prediction of thermal properties of stevia powder**

AAjit K. Mahapatra<sup>1</sup>, Ramana Gosukonda<sup>2</sup>, Bipul K. Biswas<sup>1</sup> and Hema L. Degala<sup>1</sup>. <sup>1</sup>Agricultural Research Station, <sup>2</sup>Department of Agricultural Sciences, College of Agriculture, Family Sciences and Technology, Fort Valley State University, Fort Valley, GA 31030. E-mail: [mahapatraa@fvsu.edu](mailto:mahapatraa@fvsu.edu)

Stevia leaves have been used as a natural sweetener for centuries by indigenous peoples of South America. Various research studies have shown that stevia contains ingredients that possess anti-diabetic, anti-hypertensive, anti-inflammatory, anti-cancer, and anti-diarrheal therapeutic effects. Commercially, stevia is available as an extract and in powdered form and its consumption in food and medicinal industries is increasing globally. Most of the studies on stevia are focused on the plant growing parameters, medicinal, and functional properties, whereas a very few studies have been reported on the processing and storage of stevia. Since many unit operations in the processing and storage of stevia powder involve heat transfer, it is important to understand its thermal properties. Stevia powder thermal properties data are required in the design of storage and drying equipment as well as in the estimation of process times for refrigerating, heating, or drying. An energy balance for heating or cooling of stevia powder cannot be made and the temperature profile within the powder cannot be determined without knowing the thermal properties of stevia powder. In recent years, Artificial Neural Networks (ANNs) have been used for modeling of complex and nonlinear processes. The advantages of ANNs over conventional mathematical methods in modeling performance have been recognized by many researchers. This study focused on the development of ANN models using Backpropagation and Kalman Filter learning algorithms for predicting thermal properties (thermal conductivity, volumetric specific heat, and thermal diffusivity) of stevia powder as a function of moisture content.

### **P24. Effect of medicinal plant extracts against *Fusarium oxysporum* f. sp. *albedinis*: the causal agent of Bayoud disease on date palm**

El Mehdi Bouchtaoui<sup>1</sup>, Adil Essarioui<sup>2</sup>, and Nouredine Hamamouch<sup>1</sup>. <sup>1</sup>Laboratory of Biotechnology and Plant Physiology, Faculty of Sciences, University Mohammed V, Rabat, Morocco.

<sup>2</sup>National Institute for Agricultural Research, Regional Center of Errachidia, Errachidia, Morocco.

E-mail: Nouredine Hamamouch, PhD. [hamamouch@yahoo.com](mailto:hamamouch@yahoo.com), [elmehdibouchtaoui@gmail.com](mailto:elmehdibouchtaoui@gmail.com)

*Fusarium oxysporum* f. sp. *albedinis* (Foa) is the causal agent of the vascular wilt of date palm, also known as Bayoud disease. It is the most serious soil borne disease of date palm causing the disappearance of millions of palm trees in Moroccan's oasis. Attempts to control the disease have focused on the selection of genetically resistant cultivars. However, some of the elite-but-sensitive industrial cultivars, notably "Medjool", are highly sensitive to Foa and their cultivation requires an integrated management. The objective of this study was to examine the effect of ethanolic and aqueous leaf extracts, as well as essential oils from two medical plants; *Artemisia absinthium* and *Mentha pulegium*, on Foa in-vitro. Four concentrations of plant extracts (0.625 mg/ml to 5 mg/ml) and essential oils (1 µl/ml to 7 µl/ml) were tested for their ability to inhibit spore germination, sporulation, and mycelial growth of Foa. The most significant results were obtained using the ethanolic extract of *A. absinthium* and the essential oil of *M. pulegium*. When applied at 5 mg/ml, the ethanolic extract of *A. absinthium* inhibited 100% Foa spore germination as well as sporulation, and 35% of the pathogen's mycelial growth. All other extracts showed weaker effects on Foa. In addition, the essential oil of *M. pulegium* was more effective against Foa compared to the essential oil of *A. absinthium*. The former completely inhibited Foa mycelial growth and sporulation even at a low concentration of 4 µl/ml. These results provide significant insight into the potential of using bioactive compounds from *A. absinthium* and *M. pulegium* in future management strategies of date palm Bayoud disease in Morocco.



## POSTER ABSTRACTS

### **P25. Antimicrobial and Antioxidant Properties of Central American Traditional Medicinal Plants**

Nadjet Cornejal <sup>1,2</sup>, Evian Pollack <sup>1</sup>, Rajvinder Kaur <sup>1</sup>, Ashanna Persaud <sup>1</sup>, Christine Priano <sup>1</sup>, Adolfina Koroch <sup>1</sup>, José A. Fernandez Romero <sup>1,3</sup>. <sup>1</sup>City University of New York Borough of Manhattan Community College, New York, NY 10007, <sup>2</sup>City University of New York Brooklyn College, New York, NY, 11210, <sup>3</sup>Center for Biomedical Research, Population Council, New York, NY, 10065.  
E-mail: [evian.pollack@stu.bmcc.cuny.edu](mailto:evian.pollack@stu.bmcc.cuny.edu), [jfernandezromero@bmcc.cuny.edu](mailto:jfernandezromero@bmcc.cuny.edu)

The search for alternative naturally occurring antimicrobial agents never ends, especially when emerging diseases like COVID-19 urge to find safe and effective ways to prevent or treat these infections. We studied the antioxidant properties and potential antiviral activity of three medicinal plants used in Central America: *Theobroma cacao*, *Bourreria huanita*, and *Elettaria cardamomum*. Ethanol plant extracts were prepared from commercial products derived from the seeds or flowers of these plants. The cytotoxicity and antiviral activity against severe acute respiratory syndrome coronavirus type 2 (SARS-CoV-2) were evaluated using the XTT colorimetric assay and a SARS-CoV-2 delta pseudoviral model, respectively. The half-maximal cytotoxic concentration (CC<sub>50</sub>) and half-maximal effective concentration (EC<sub>50</sub>) were used to calculate the therapeutic index (TI). We estimated total phenolics and antioxidant activity using Sigma Aldrich's phenolic compounds and total antioxidant assay kits. The extracts showed total phenolics in the range of 0.23 to 1.36 mM catechin equivalents, with *T. cacao* beans extract showing the highest content. The antioxidant activity showed values between 0.05 to 0.34 mM Trolox equivalents, and once again, *T. cacao* beans extract showed the highest antioxidant activity. Most plant extracts showed none to moderate selective antiviral activity with TI values between 1 and 5.5. However, *T. cacao* beans extract had excellent antiviral activity against SARS-CoV-2 with a CC<sub>50</sub> and EC<sub>50</sub> of 425.1 and 19.7 µg/mL, respectively, resulting in a TI value of 22. Future studies will look at the phytochemical composition of this extract to identify potential active pharmaceutical ingredients.

---

### **P26. The experimental evaluation of *Chlorella vulgaris* extracellular polysaccharide anti-asthmatic profile**

Martina Sutovska <sup>1</sup>, Peter Capek <sup>2</sup>, Jana Barborikova <sup>1</sup> and Sona Franova <sup>1</sup>. <sup>1</sup>Department of Pharmacology Jessenius School of Medicine, Comenius University, Martin, Slovakia; <sup>2</sup>Institute of Chemistry, Center of Glycomics, Slovak Academy of Sciences, Bratislava, Slovakia.  
E-mail: [martina.sutovska@uniba.sk](mailto:martina.sutovska@uniba.sk)

Cultivation of green microalgal species *Chlorella vulgaris* resulted a significant production of extracellular polysaccharide (EPS). The chemico-spectroscopic studies on EPS revealed its molecular profile and very complex highly branched  $\alpha$ -L-arabino- $\alpha$ -L-rhamno- $\alpha,\beta$ -D-galactan structure. Biological activity tests showed that EPS caused significant antitussive effect (on chemically-induced cough reflex), bronchodilatory activity (on airway reactivity elicited by inhalation of histamine and methacholine) and anti-inflammatory effect (evidenced using Bioplex® assay) in guinea pigs with experimentally induced allergic airway inflammation. Precautionary administrated of EPS suppresses both, airway hyperreactivity and airway inflammation. Bronchodilation was similar to the effect of a classic anti-asthmatic bronchodilator - salmeterol and suppression of allergic inflammation was comparable with that of budesonide, an anti-inflammatory corticosteroid agent. The combination of *Chlorella* EPS with budesonide possessed synergic calming effect on airway hyperreactivity and prevented cilio-inhibitory effect of this biopolymer. Therefore, *chlorella* EPS appears to be a promising agent for the prevention of chronic airway inflammation, which is the basic pathogenetic mechanism of many respiratory diseases, including bronchial asthma. Ex vivo biological assays showed EPS-induced increase in interleukin-12 (IL-12) and interferon-gamma (INF- $\gamma$ ) levels corresponding to transforming growth factor beta (TGF- $\beta$ ) decrease in experimental guinea pig asthma model. These facts point to the anti-remodeling effect of *chlorella* EPS and confirm its possible application in the treatment of asthma and chronic obstructive pulmonary disorder.

## POSTER ABSTRACTS

### P27. In vitro antiviral activity of essential oils against SARS-CoV-2 Delta variant

Luiz Torres Neto <sup>1,2,3</sup>, Maria Lucia Guerra Monteiro <sup>1,2,3</sup>, José Fernandez-Romero <sup>4,5</sup>, Natalia Teleshova <sup>5</sup>, James Sailer <sup>5</sup> and Carlos Adam Conte Junior <sup>1,2,3</sup>. <sup>1</sup> COVID-19 Research Group, Center for Food Analysis (NAL), Technological Development Support Laboratory (LADETEC), Cidade Universitária, Rio de Janeiro 21941-598, Brazil, <sup>2</sup> Laboratory of Advanced Analysis in Biochemistry and Molecular Biology (LAABBM), Department of Biochemistry, Federal University of Rio de Janeiro (UFRJ), Cidade Universitária, Rio de Janeiro 21941-909, Brazil, <sup>3</sup> Graduate Program in Food Science (PPGCAL), Institute of Chemistry (IQ), Federal University of Rio de Janeiro (UFRJ), Cidade Universitária, Rio de Janeiro 21941-909, Brazil, <sup>4</sup> Science Department, Borough of Manhattan Community College, The City University of New York, NY, 199 Chambers Street, Science Department room N699, New York, NY 10007, USA. <sup>5</sup> Center for Biomedical Research, Population Council, 1230 York Avenue, New York, NY 10065, USA.

E-mail: [luiz-torres-neto@hotmail.com](mailto:luiz-torres-neto@hotmail.com); [marialuciaguerra@yahoo.com.br](mailto:marialuciaguerra@yahoo.com.br); [jromero@popcouncil.org](mailto:jromero@popcouncil.org); [nteleshova@popcouncil.org](mailto:nteleshova@popcouncil.org); [jsailer@popcouncil.org](mailto:jsailer@popcouncil.org); [conte@iq.ufrj.br](mailto:conte@iq.ufrj.br)

Essential oils (EOs) are widely studied due to their broad bioactive spectrum, including antimicrobial, antifungal, antioxidant, and antiviral properties. These properties are attributed to terpenes present in their composition. Recent studies have suggested the antiviral potential of EOs against SARS-CoV-2 due to their proven activity against several other viruses (e.g., herpes simplex, influenza, human immunodeficiency virus, among others) and for managing symptoms in patients with COVID-19. This study aimed to evaluate the in vitro antiviral action of 12 EOs against the SARS-CoV-2 Delta variant using a pseudoviral (PsV) model, which is rare in the literature. The twelve EOs assessed in this study were selected based on our preliminary systematic and meta-analytic study to identify the most promising EOs against SARS-CoV-2 considering their affinity by target proteins from in silico studies. Furthermore, EOs with antiviral potential against other viruses identified from the literature were also selected. Antiviral activity and cytotoxicity were tested using the XTT and cell-based PsV assays, and the half-maximal effective concentration (EC<sub>50</sub>) and half-maximal cytotoxic concentration (CC<sub>50</sub>) was determined (GraphPad Prism software version 9.0.2). The therapeutic index (TI) was calculated for each EO (TI = CC<sub>50</sub>/EC<sub>50</sub>). The EC<sub>50</sub> values ranged from 0.09 to 57 µg/mL, wherein the EOs from *Syzygium aromaticum* (0.09 µg/mL), *Cymbopogon citratus* (0.1 µg/mL), *Citrus limon* (0.15 µg/mL), *Pelargonium graveolens* (0.2 µg/mL), *Origanum vulgare* (0.4 µg/mL) and *Illicium verum* (0.5 µg/mL) were more effective against the virus. The CC<sub>50</sub> values ranged from 0.2 to 81 µg/mL, resulting in the highest TI values for *Illicium verum* (TI = 60), *Citrus limon* (TI = 8.67), *Pelargonium graveolens* (TI = 8.50), and *Syzygium aromaticum* (TI = 4.44). These four EOs proved to be a promising EO for SARS-CoV-2 inhibition, mainly *Illicium verum*.



## POSTER ABSTRACTS

### **P28. Evaluating the Potential Antiproliferative Effects of *Ulva lactuca* in Colorectal Cancer Cell Lines HT-29 and HCT-116**

Rocío del Mar Avilés-Mercado<sup>1</sup>, Ángel R. Núñez-Marrero<sup>1</sup>, Rosalinda Aybar-Batista<sup>1</sup>, Jacobo Morales-Méndez<sup>1</sup>, Alok Arun<sup>1</sup>, Juan A. Negrón-Berríos<sup>1</sup>. <sup>1</sup>Institute of Sustainable Biotechnology (ISBiot), Inter American University of Puerto Rico-Barranquitas Campus, Puerto Rico, USA 00794. E-mail: rocavi7173@br.uipr.edu

Colorectal cancer (CC) is one of the three most frequent cancers in Puerto Rico. It is also one of the top three causes of cancer deaths. The traditional treatments for CC although moderately effective, cause detrimental side effects to patients. For this reason, the search for alternative cures that can be both efficient and less harmful is highly desirable. Marine macroalgae contain different compounds with important medical properties such as antimicrobial, antioxidant, antitumor, and antiviral. Green macroalgae such as *Ulva* has been described as having potential anti-cancer properties. However, there are few studies devoted to evaluating such algae anticancer properties in the context of CC. The aim of this research was to evaluate the potential effects of the *Ulva lactuca* in the proliferation of the metastatic colorectal cancer cell lines HT-29 and HCT-116. These were cultured and exposed for 24 hours, 48 hours, and 72 hours to methanolic extracts made of *U. lactuca* from Isla Magueyes (southwest area of Puerto Rico). Subsequently, a MTT assay was performed to evaluate the proliferation and viability of both cell lines. Our results highlighted a reduction of the proliferation and cell viability of both cell lines to specific dilutions of the methanolic extracts. Such results prompt further analyses regarding the potential mechanisms by which *U. lactuca* affects metastatic CC cell lines. This would represent a novel aspect with the potential of improving treatment options against a highly frequent and aggressive cancer type whose current treatments options are not suitable for everybody. The funding for this research was provided by NSF HSI-STEM Award No. 1928792.

---

### **P29. Discovery and development of anxiolytic agents from tropical marine macroalgae**

Ricardo M. Cruz-Sánchez<sup>1</sup>, Gabriela M. Ramírez-Renta<sup>1</sup>, Geraldine M. Ortiz-Sosa<sup>2</sup>, Aniel J. Rivera-Arzola<sup>3</sup>, Janiennid Alicea-Tirado<sup>3</sup>, Adrián E. González-Santiago<sup>2</sup>, Grayce E. Dyer Levin<sup>4</sup>, Eduardo Caro-Díaz<sup>4</sup>, Ricardo Chiesa<sup>1</sup>. <sup>1</sup>Department of Biology, <sup>2</sup>Natural Sciences General Program, <sup>3</sup>Department of Chemistry, University of Puerto Rico, Cayey, Puerto Rico, and <sup>4</sup>UPR – Medical Science Campus. E-mail: ricardo.cruz18@upr.edu; ricardo.chiesa@upr.edu

Surveys show that one-third of the population is affected by an anxiety disorder. These are mostly treated with benzodiazepines (BZD's) which can induce dependence disorders through tolerance and resistance mechanisms. This highlights the need for safer and more effective anxiolytic drugs. Recently, marine natural products have proven to be a rich chemical space for drug discovery and development. Specifically, tropical marine algae produce a wide range of metabolites with diverse biological activity, including neuroprotection. We propose to study the anxiolytic effects of natural products derived from marine tropical brown algae using an invertebrate fly model (*Drosophila*). Our research aims to chemically characterize brown algae extracts, use these extracts for anxiety-related behavioral tests using *Drosophila* and identify the chemical components responsible for the anxiolytic activity. We have recently published results reporting that a crude extract of *Stypopodium zonale* decreases fear-related behavior in *Drosophila melanogaster* using an Open Field Test (OFT). This study will further develop and validate *Drosophila* as a versatile and useful alternative for drug discovery efforts geared towards the treatment of anxiety disorders. Our research represents an unprecedented approach to anxiolytic drug discovery. It will also contribute to our understanding of Puerto Rico's marine algae and their natural product's chemo-diversity. Our marine brown algae chemical library will also serve as a unique resource that harnesses Puerto Rico's marine biodiversity and platform for other biotechnological discoveries.



## POSTER ABSTRACTS

### **P30. Fermentation of *Vaccinium floribundum* Berries Reduces Oxidative Stress in Endothelial Cells**

Luisa Marracino<sup>1</sup>, Angela Punzo<sup>3</sup>, Paolo Severi<sup>1</sup>, Achille Wilfred Ouambo Talla<sup>1</sup>, Alessia Ascierto<sup>1</sup>, Paola Rizzo<sup>1,2</sup>, Cristiana Caliceti<sup>4,5,6</sup> <sup>1</sup>Department of Translational Medicine and Laboratory for Technologies of Advanced Therapies (LTTA), University of Ferrara, 44121 Ferrara, Italy. <sup>2</sup>Maria Cecilia Hospital, GVM Care & Research, 48033 Cotignola, Italy. <sup>3</sup>Department of Chemistry "Giacomo Ciamician" Alma Mater Studiorum, University of Bologna, 40126 Bologna, Italy.

<sup>4</sup>Biostructures and Biosystems National Institute (INBB), 00136 Rome, Italy. <sup>5</sup>Department of Biomedical and Neuromotor Sciences, Alma Mater Studiorum, University of Bologna, 40126 Bologna, Italy. <sup>6</sup>Interdepartmental Centre for Renewable Sources, Environment, Sea and Energy (CIRI FRAME), Alma Mater Studiorum, University of Bologna, 40126 Bologna, Italy.

E-mail: rzzpla@unife.it, alessia.asciarto@edu.unife.it, achillewilfred.ouambotalla@unife.it, mrrlsu@unife.it, svrpla@unife.it, angela.punzo2@unibo.it, cristiana.caliceti@unibo.it

Evidence suggests that high consumption of food containing antioxidants reduces the risk of cardiovascular diseases. Additionally, fruits fermentation with lactic acid bacteria (LAB) enhances their beneficial properties as regulators of the immune, digestive, and cardiovascular system. In this context, we investigated the effects of fermentation with *Lactiplantibacillus plantarum* of Pushgay berries (*Vaccinium floribundum*, Ericaceae family) in human umbilical vein endothelial cells (HUVECs). The effects of berries solutions on cell viability and proliferation were assessed by Lactate dehydrogenase (LDH) release and Trypan blue exclusion test. Antioxidant activity was evaluated by a cell-based chemiluminescent probe for the detection of intracellular H<sub>2</sub>O<sub>2</sub> production. Heme oxygenase-1 (HO-1) expression levels were investigated by RT-qPCR. Glutathione reductase (GR), glutathione peroxidase (Gpx), superoxide dismutase (SOD), and catalase (CAT) activities, as markers of intracellular antioxidant defense, were evaluated by spectrophotometric analysis. Data showed that fermentation of Pushgay berries increases their intracellular antioxidant activity, as indicated by the reduction in H<sub>2</sub>O<sub>2</sub>-induced cell death and the decrease in H<sub>2</sub>O<sub>2</sub>-induced HO-1 gene expression in HUVECs treated for 24 h with fermented berries solution (10 µg/mL). Taken together, our results show that LAB fermentation of Pushgay berries enhances their antioxidant properties.



## POSTER ABSTRACTS

### **P31. The potential application of essential oil of *Thymus broussonetii* Boiss., as a preservative ingredient in cosmetic products**

Smahane Boukhira<sup>1</sup>, Nouredine Hamamouch<sup>2</sup>, Dalila Boustia<sup>1</sup>. <sup>1</sup>Laboratory of Biotechnology, Environment, Agrifood, and Health, Faculty of Sciences, University Sidi Mohamed Ben Abdellah, Fez 30050, Morocco. <sup>2</sup>Laboratory of Biotechnology and Plant Physiology, Faculty of Sciences, University Mohammed V, Rabat, Morocco; E-mail: smahaneboukhira@gmail.com ; hamamouch@yahoo.com

The aim of this study was to evaluate the antioxidant, and preservative efficacy of *Thymus broussonetii* Boiss. essential oil (EO) in topically applied formulation using a challenge test. The EO was extracted from the aerial part of *T. broussonetii* using hydrodistillation and the obtained EO was further analysed by GC/MS (Gas Chromatography/Mass Spectrometry). The antioxidant effect of the EO was evaluated using two methods, the inhibition of free radical 2,2-DiPhenyl-1-PicrylHydrazyl (DPPH) and Reducing Power Ferric Antioxidant Power (FRAP) methods. Furthermore, the preservative efficacy of *T. broussonetii* EO was assayed at 2% (v/w) in topical cream formulation, using a challenge test against standard specific microorganisms recommended by the European Pharmacopoeia. Chemical composition of the EO revealed the presence of thymol (63.09%), p-cymene (11%) and  $\gamma$ -terpinene (8.99%) as the major components. The antioxidant assays revealed that the essential oil exhibited strong antioxidant activity as indicated by the minimum inhibitory concentration IC<sub>50</sub> (IC<sub>50</sub> = 210 $\pm$ 0.3  $\mu$ g/mL for DPPH assay and IC<sub>50</sub> = 84 $\pm$ 0.21  $\mu$ g/mL for FRAP assay) when compared to Quercetin and Butylatedhydroxytoluene (BHT) as controls. Moreover, the EO of *T. broussonetii* evaluated at 2% (v/w) in cream formulation succeeded in satisfying the A criteria for preservation efficacy against *Staphylococcus aureus*, *Escherichia coli* and *Aspergillus brasiliensis* but exhibited less efficacy against *Pseudomonas aeruginosa* (1.78 log reduction in the number of CFU/g after the 7 day of evaluation) and *Candida albicans* (1.09 log reduction in the number of CFU/g after the 14 days of evaluation) when compared to the synthetic preservative Phenoxyethanol 1% (v/w). These results indicate that the EO of *T. broussonetii* possess an important antimicrobial and antioxidant activities and can therefore be used as a natural preservative ingredient in cosmetic industry.



# 11<sup>th</sup> ANNUAL HYBRID CONFERENCE

**American Council for Medicinally Active Plants (ACMAP)**

EXPLORING THE UNEXPLORED: SCIENCE AND APPLICATIONS OF MEDICINAL PLANTS

JUNE 28 - JULY 2, 2022 | PUERTO RICO, USA

## SPONSORS





**11<sup>th</sup> ANNUAL HYBRID CONFERENCE**  
**American Council for Medicinally Active Plants (ACMAP)**  
**EXPLORING THE UNEXPLORED: SCIENCE AND APPLICATIONS OF MEDICINAL PLANTS**  
**JUNE 28 - JULY 2, 2022 | PUERTO RICO, USA**

# INSTITUTE OF SUSTAINABLE BIOTECHNOLOGY

The Institute of Sustainable Biotechnology (ISBiot) at Barranquitas Campus of Inter American University of Puerto Rico is dedicated to promoting and educating the public in the area of modern biotechnology. As an organization, we are committed to fostering the development of professionals who contribute to the progress of Biotechnology as a key piece for the evolution and socio-economic development of our society. The commitment of the Institute of Sustainable Biotechnology is to link the knowledge and strategies of Biotechnology with sustainable practices to promote learning, research and development in Puerto Rico. The following specialized courses are currently being offered at the graduate level in ISBiot:



## **Master of Science in Biotechnology**

The Master of Science in Biotechnology with Specializations in Biotechnology of Plants and Industrial Biotechnology has as its goal the formation of professionals who possess the knowledge, skills, attitudes and the social commitment, which allow them to make responsible contributions to the scientific development for our society and to the advancement of biotechnology. It aims to facilitate the development of graduates that can handle the basic technologies of biosciences and emit judgments related to their area of specialization and to related areas.

## **Doctor of Philosophy (PhD) in International Biotechnology**

The Doctoral Program in International Biotechnology, within its general goals, aims to train professionals focused on mastering the knowledge, skills and attitudes required in the area of biotechnology in an international context. The program concentrates its interest in scientific research and technological innovation in agriculture, industry, the environment, and health. The PhD seeks to contribute to the formation of values and attitudes, within the ethical, legal and social framework that should characterize the professional in the field of Biotechnology. In addition, it recognizes the importance of scientific research in solving real problems that affect the quality of life of societies globally. The profile of academic competences will allow to find solutions to problems from a critical, creative and ethical perspective.



For eligibility requirements, please contact:

### **Institute of Sustainable Biotechnology**

Department of Science and Technology  
Inter American University of Puerto Rico  
Barranquitas Campus, PO Box 517  
Barranquitas, Puerto Rico 00794

Telephone: 787-857-3600

Ext. #2067 | #2089

Webpage: [www.br.inter.edu](http://www.br.inter.edu)





# **11<sup>th</sup> ANNUAL HYBRID CONFERENCE**

## **American Council for Medicinally Active Plants (ACMAP)**

**EXPLORING THE UNEXPLORED: SCIENCE AND APPLICATIONS OF MEDICINAL PLANTS**

**JUNE 28 - JULY 2, 2022 | PUERTO RICO, USA**

### **ACMAP INCOMING BOARD OF DIRECTORS (8/1/22 – 7/31/23)**

Executive Director: Fabricio Medina-Bolivar, Ph.D. (Arkansas State University, USA)

President: Nirmal Joshee, Ph.D. (Fort Valley State University, Georgia)

Vice President: Joshua Kellogg, Ph.D. (Pennsylvania State University, Pennsylvania)

Secretary: Andrea Doseff, Ph.D. (Michigan State University, Michigan)

Treasurer: Sun-Ok Lee, Ph.D. (University of Arkansas, Arkansas)

Member-at-Large (1): Alok Arun, Ph.D. (Inter American University of Puerto Rico, Puerto Rico)

Member-at-Large (2): Chhandak Basu, Ph.D. (California State University, California)

Member-at-Large (3): Jeremy Johnson, Ph.D. (University of Illinois at Chicago, Illinois)

ACMAP Ambassador (1) Samantha Sherman (Fort Valley State University, Georgia)

ACMAP Ambassador (2) Emily Merchant (Rutgers University, New Jersey)

### **ACMAP CURRENT BOARD OF DIRECTORS**

Executive Director: Fabricio Medina-Bolivar, Ph.D. (Arkansas State University, USA)

President: Anait S. Levenson, M.D., Ph.D. (Long Island University, New York)

Vice President: Joshua Kellogg, Ph.D. (Pennsylvania State University, Pennsylvania)

Secretary: Chhandak Basu, Ph.D. (California State University, California)

Treasurer: Sun-Ok Lee, Ph.D. (University of Arkansas, Arkansas)

Member-at-Large (1): Alok Arun, Ph.D. (Inter American University of Puerto Rico, Puerto Rico)

Member-at-Large (2): Andrea Doseff, Ph.D. (Michigan State University, Michigan)

Member-at-Large (3): Nirmal Joshee, Ph.D. (Fort Valley State University, Georgia)





# **11<sup>th</sup> ANNUAL HYBRID CONFERENCE**

## **American Council for Medicinally Active Plants (ACMAP)**

**EXPLORING THE UNEXPLORED: SCIENCE AND APPLICATIONS OF MEDICINAL PLANTS**

**JUNE 28 - JULY 2, 2022 | PUERTO RICO, USA**

### **ACMAP 2022 COMMITTEE MEMBERS**

#### **Inter American University of Puerto Rico Barranquitas Campus**

Abner I. Hernández González  
Aixa Serrano Febo  
Alok Arun, Ph.D.  
Carmen C. Rosado Berríos  
Cristian J. Ríos Colón  
Denisse Meléndez Serrano  
Eleane Rosado López  
Jacobó Morales Méndez  
Jonathan Ortiz Morales  
Joseph Ortiz Arroyo  
María M. Meléndez Ortega  
Michelle Cartagena Rivera, Ph.D.  
Rita Alvarado Rojas  
Rosalinda Aybar Batista  
Rosenid Vega Colón  
Sonia I. Burgos Rivera  
Zullymar Ortiz Rivera

#### **Inter American University of Puerto Rico Bayamón Campus**

Dr. Rafael Canales  
Ángel Martínez  
Herinaldo Torres





# **11<sup>th</sup> ANNUAL HYBRID CONFERENCE**

## **American Council for Medicinally Active Plants (ACMAP)**

**EXPLORING THE UNEXPLORED: SCIENCE  
AND APPLICATIONS OF MEDICINAL PLANTS**

**JUNE 28 - JULY 2, 2022 | PUERTO RICO, USA**

**Inter American University of Puerto Rico  
Barranquitas Campus**

P.O. Box 517  
Barranquitas, P.R. 00794  
Telephone: 787-857-3600  
[www.br.inter.edu](http://www.br.inter.edu)

**Inter American University of Puerto Rico  
Bayamón Campus**

500 Dr. John Will Harris Street  
Bayamón, 00957  
Telephone: (787) 279-1912  
<https://interbayamon3.azurewebsites.net/>

**Copyright © 2022 All rights reserved.**